For Research Use Only

VIP Polyclonal antibody

Catalog Number: 16233-1-AP

4 Publications



Basic Information

Catalog Number: 16233-1-AP

Source:

GenBank Accession Number: BC009794

Purification Method: Antigen affinity purification

GeneID (NCBI): Size: 400 μg/ml

7432 **UNIPROT ID:**

Rabbit P01282 Isotype: Full Name:

vasoactive intestinal peptide

Calculated MW: Immunogen Catalog Number: AG8933 169 aa, 19 kDa

Positive Controls:

IHC: human colon tissue, rat small intestine tissue, mouse brain tissue, human small intestine tissue

Recommended Dilutions:

IHC 1:400-1:1600 IF 1:50-1:500

IF: mouse colon tissue, COLO 320 cells

Applications

Tested Applications: IF/ICC,IF-P, IHC, ELISA Cited Applications: WB, IF, IHC

Species Specificity: human, mouse, rat Cited Species: rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Vasoactive intestinal peptide (VIP), a short peptide containing 28 amino acids belonging to the secretin-glucagon family, is initially isolated from the gastrointestinal tract as a potent vasodilator peptide. VIP was initially identified in normal nervous tissue and neurons and was subsequently recognized as a neurotransmitter widely distributed in various tissues. The wide distribution of VIP determines its involvement in a range of biological activities, such as gut motility, hormonal regulation, circadian rhythms, immune responses, and carcinogenesis. The general physiologic effects of VIP include vasodilation, anti-inflammatory actions, cell proliferation, hormonal secretion, regulation of gastric motility, and smooth muscle relaxation; therefore, VIP has emerged as a promising drug candidate for the treatment of several diseases.

Notable Publications

Author	Pubmed ID	Journal	Application
Muhua Huang	34493162	Pharm Biol	IF
Marta Gil	38540797	Biomolecules	WB
Yixi Li	37506142	Immun Inflamm Dis	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16233-1-AP (VIP antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16233-1-AP (VIP antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using VIP antibody (16233-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Calretinin antibody (66496-1-Ig, Clone: 2D7A9, red).



Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using VIP antibody (16233-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Calretinin antibody (66496-1-Ig, Clone: 2D7A9, red).