

OCIAD1 Polyclonal antibody

 Catalog Number: 16634-1-AP 2 Publications

Basic Information

Catalog Number: 16634-1-AP	GenBank Accession Number: BC003409	Purification Method: Antigen affinity purification
Size: 350 µg/ml	GeneID (NCBI): 54940	Recommended Dilutions: WB 1:500-1:2000 IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF/ICC 1:10-1:100
Source: Rabbit	UNIPROT ID: Q9NX40	
Isotype: IgG	Full Name: OCIA domain containing 1	
Immunogen Catalog Number: AG9977	Calculated MW: 28 kDa	
	Observed MW: 29-35 kDa	

Applications

Tested Applications: IF/ICC, IHC, IP, WB, ELISA	Positive Controls: WB : HeLa cells, MCF-7 cells, human placenta tissue IP : HeLa cells, IHC : human liver cancer tissue, human kidney tissue, human pancreas cancer tissue, human thyroid cancer tissue IF/ICC : HepG2 cells,
Cited Applications: WB, IF, IHC	
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

OCIAD1 was first identified by immunoscreening of an ovarian carcinoma cDNA expression library with ascites fluid from ovarian cancer patients (PMID: 11162530). OCIAD1 has been reported as a key player in ovarian cancer cell adhesion, as well as a key player in generating ovarian cancer recurrence (PMID: 18328549; 20515946). In addition to its roles in cancer, OCIAD1 participates in maintaining stem cell potency by regulating the Jak/STAT pathway (PMID: 23972987). Several alternatively spliced forms of OCIAD1 gene have been identified. The longest form (1.4 kb) is predicted to encode for a 27.6 kDa protein of 245 amino acids. This antibody detects OCIAD1 with an apparent molecular weight of ~35 kDa as has been demonstrated by several researches (PMID: 27345969; 27345976).

Notable Publications

Author	Pubmed ID	Journal	Application
Huong T L Tran	32697788	PLoS One	WB,IF
Nagata Chigusa C	22726067	Pathol Int	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16634-1-AP (OCIAD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 16634-1-AP (OCIAD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 16634-1-AP (OCIAD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells, using OCIAD1 antibody 16634-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-OCIAD1 (IP:16634-1-AP, 4ug; Detection:16634-1-AP 1:300) with HeLa cells lysate 3200ug.