# For Research Use Only

# RPS13 Polyclonal antibody

Catalog Number: 16680-1-AP

4 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 16680-1-AP BC006772

Size: GeneID (NCBI): 6207

Source: UNIPROT ID: Rabbit P62277

IgG ribosomal protein S13
Immunogen Catalog Number: Calculated MW:
AG10008 17 kDa

Observed MW: 14 kDa, 17 kDa

Full Name:

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:1000-1:4000 IHC 1:50-1:500 IF 1:50-1:500

**Applications** 

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications:

WB

Isotype:

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls

WB: MCF-7 cells, mouse oterus tissue, rat oterus tissue

IHC: human colon cancer tissue,

IF: HeLa cells,

# **Background Information**

RPS13, also named as 40S ribosomal protein S13, belongs to the ribosomal protein S15P family. Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. RPS13 is located in the cytoplasm. It is co-transcribed with two U14 small nucleolar RNA genes, which are located in its third and fifth introns. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Keely Oltion	36638793	Cell	WB
Yanrong Zhuang	37369203	Cell	WB
Peng Shuping S	21425412	Stem Cells	WB

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 16680-1-AP (RPS13 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using 16680-1-AP (RP513 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 16680-1-AP (RPS13 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).