For Research Use Only

GFAP Polyclonal antibody, PBS Only

Catalog Number:16825-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

16825-1-PBS

BC013596 GeneID (NCBI): 2670

1 mg/ml Source: Rabbit

Size:

UNIPROT ID: P14136 Full Name:

Isotype:

glial fibrillary acidic protein

Immunogen Catalog Number: AG10423

Calculated MW: 432 aa, 50 kDa

Observed MW:

45-50 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, IF-Fro, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

GFAP (Glial fibrillary acidic protein), an intermediate-filament (IF) protein, is specifically expressed in cells of astroglial lineage and is widely used to mark astroglia in the brain. It is also used as a marker for intracranial and intraspinal tumors arising from astrocytes.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80 $^{\circ}\text{C}$

Storage Buffer: PBS Only


Selected Validation Data




Immunohistochemical analysis of paraffinembedded Hippocampus tissue slide from Alzheimer Mouse Model using 16825-1-AP (GFAP antibody) at dilution of 1:50000 (under 10x lens). Data from NeuroScience Associates, Inc. This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using GFAP antibody (16825-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using 16825-1-AP (GFAP antibody) at dilution of 1:1000 and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using GFAP antibody (16825-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16825-1-AP (GFAP antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16825-1-AP (GFAP antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.




WB result of GFAP antibody (16825-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GFAP transfected U-251 cells. This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 16825-1-AP (GFAP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using GFAP antibody (16825-1-AP) at dilution of 1:200 and Coralite®594-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-4). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed A172 cells using GFAP antibody (16825-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 16825-1-PBS in a different storage buffer formulation.