For Research Use Only

YTHDF1 Polyclonal antibody Catalog Number: 17479-1-AP Featured Product 264


Featured Product 264 Publications




Basic Information	Catalog Number: 17479-1-AP	GenBank Accession Number: BC050284		Purification Method: Antigen affinity purification	
	Concentration: GeneID (NCBI):):	Recommended Dilutions: WB: 1:1000-1:8000	
	800 ug/ml	54915			
	Source: UNIPROT ID: Rabbit Q9BYJ9 Isotype: Full Name:		IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate		
				IHC: 1:250-1:1000 IF-P: 1:200-1:800	
	lgG	YTH domain fa	amily, member 1	IF/ICC: 1:200-1:800	
	Immunogen Catalog Number: AG11509	Calculated MV 559 aa, 61 kDa			
		Observed MW: 60 kDa			
Applications			Positive Co	ntrols:	
	WB, IHC, IF/ICC, IF-P, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP, RIP		tissue, rat b	WB : HeLa cells, HepG2 cells, A549 cells, mouse brai tissue, rat brain tissue, HEK-293 cells, Jurkat cells, MCF-7 cells, NIH/3T3 cells	
	Species Specificity:		IP : mouse l	IP : mouse brain tissue,	
			IHC : huma	n kidney tissue, human hypothalamus	
	human, mouse, rat, pig, monkey, chicken, zebrafish		tissue		
	Note-IHC: suggested antigen retrieval with			IF-P: mouse brain tissue,	
			IF/ICC : U2	US cells,	
Background Information	YTHDF1, also named YTH domain-containing family protein 1 or C20orf21, is a 559 amino acid protein, which localizes in the cytoplasm. YTHDF1 specifically recognizes and binds N6-methyladenosine (m6A)-containing mRNAs, and promotes mRNA translation efficiency. M6A is a modification present at the internal sites of mRNAs ar some non-coding RNAs and plays a role in the efficiency of mRNA splicing, processing, and stability. YTHDF1 acts a a regulator of mRNA translation efficiency: promotes ribosome loading to m6A-containing mRNAs and interacts wi translation initiation factors eIF3 (EIF3A or EIF3B) to facilitate translation initiation. YTHDF1 exists two isoforms and the calculated molecular weight of isoforms are 61 kDa and 21 kDa.				
Notable Publications	Author	Pubmed ID	Journal	Application	
Notable Publications	Fanpeng Zhao	34593014	Mol Neurodegener		
Notable Publications			1 Dial Cham	14/D	
Notable Publications	Kusuma Suphakhong	36183833	J Biol Chem	WB	
Notable Publications	Kusuma Suphakhong Bin Zhang	36183833 32964498	Neurochem Res	WB	

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of YTHDF1 antibody (17479-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-YTHDF1 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 17479-1-AP (YTHDF1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




A549 cells were subjected to SDS PAGE followed by western blot with 17479-1-AP (YTHDF1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using YTHDF1 antibody (17479-1-AP) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 17479-1-AP (YTHDF1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 17479-1-AP (YTHDF1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 17479-1-AP (YTHDF1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-YTHDF1 (IP:17479-1-AP, 4ug; Detection:17479-1-AP 1:1000) with mouse brain tissue lysate 2640ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using YTHDF1 antibody (17479-1-AP) at dilution of 1:400 and Multi-Ab Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).