For Research Use Only

MTA2 Polyclonal antibody

Catalog Number: 17554-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17554-1-AP BC053650 GeneID (NCBI): Size: 350 μg/ml 9219 **UNIPROT ID:** Source: Rabbit 094776 Full Name: Isotype:

metastasis associated 1 family,

AG11713 Calculated MW: 668 aa, 75 kDa Observed MW:

70-75 kDa

member 2

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, CoIP, ChIP Species Specificity: human, mouse, rat

Immunogen Catalog Number:

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, Jurkat cells, mouse thymus tissue,

Purification Method:

WB 1:2000-1:10000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

HEK-293 cells

IP: mouse thymus tissue,

IHC: human colon tissue, human heart tissue

Background Information

Metastasis-associated protein MTA2, also named as MTA1L1 or PID, is a 668 amino acid protein, which contains one BAH domain, one ELM2 domain, one GATA-type zinc finger and one SANT domain. MTA2 localizes in the nucleus and is widely expressed in many tissues. MTA2 may be involved in the regulation of gene expression as repressor and activator.

Notable Publications

Author	Pubmed ID	Journal	Application
Orhi Barroso-Gomila	34795231	Nat Commun	WB
Su-Li Dai	33340431	Cancer Sci	ColP,ChIP
Dapeng Wang	35201651	EMBO Rep	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of MTA2 antibody (17554-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MTA2 transfected HEK-293 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 17554-1-AP (MTA2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon using 17554-1-AP (MTA2 antibody) at dilution of 1:100 (under 40x lens).



Immunohistochemical analysis of paraffinembedded human colon using 17554-1-AP (MTA2 antibody) at dilution of 1:100 (under 10x lens).



IP result of anti-MTA2 (IP:17554-1-AP, 4ug; Detection:17554-1-AP 1:500) with mouse thymus tissue lysate 2800ug.