## For Research Use Only

# MTA2 Polyclonal antibody

Catalog Number: 17554-1-AP

Featured Product

5 Publications



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 17554-1-AP
 BC053650

 Size:
 GeneID (NCBI):

 350 ug/ml
 9219

 Source:
 UNIPROT ID:

Rabbit 094776 Isotype: Full Name:

lgG metastasis associated 1 family,
member 2

AG11713 Calculated MW: 668 aa, 75 kDa

Observed MW: 70-75 kDa Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:500-1:2000

**Applications** 

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, CoIP, ChIP Species Specificity:

Immunogen Catalog Number:

human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: HeLa cells, Jurkat cells, mouse thymus tissue,

HEK-293 cells

IP: mouse thymus tissue,

IHC: rat lung tissue, human heart tissue, rat colon

issue

# **Background Information**

Metastasis-associated protein MTA2, also named as MTA1L1 or PID, is a 668 amino acid protein, which contains one BAH domain, one ELM2 domain, one GATA-type zinc finger and one SANT domain. MTA2 localizes in the nucleus and is widely expressed in many tissues. MTA2 may be involved in the regulation of gene expression as repressor and activator.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Orhi Barroso-Gomila	34795231	Nat Commun	WB
Su-Li Dai	33340431	Cancer Sci	ColP,ChIP
Dapeng Wang	35201651	EMBO Rep	WB

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**



WB result of MTA2 antibody (17554-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MTA2 transfected HEK-293 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 17554-1-AP (MTA2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-MTA2 (IP:17554-1-AP, 4ug; Detection:17554-1-AP 1:500) with mouse thymus tissue lysate 2800ug.

Immunohistochemical analysis of paraffinembedded rat colon tissue slide using 17554-1-AP (MTA2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat lung tissue slide using 17554-1-AP (MTA2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).