### For Research Use Only

# HMGCS1 Polyclonal antibody

Catalog Number: 17643-1-AP

Featured Product

35 Publications



**Basic Information** 

Catalog Number: 17643-1-AP Size:

700 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG11825

GenBank Accession Number:

BC083514
GeneID (NCBI):
3157
UNIPROT ID:
Q01581
Full Name:

3-hydroxy-3-methylglutaryl-Coenzyme A synthase 1 (soluble)

Calculated MW: 520 aa, 57 kDa Observed MW:

57 kDa

Positive Controls:

WB: A431 cells, A549 cells, HepG2 cells, LNCaP cells,

**Purification Method:** 

WB 1:500-1:1000

IF/ICC 1:400-1:1600

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

mouse liver tissue

IP: mouse liver tissue,

IF/ICC: HepG2 cells,

**Applications** 

WB, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF, CoIP Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

**Tested Applications:** 

# **Background Information**

HMGCS1(Hydroxymethylglutaryl-CoA synthase, cytoplasmic) is also named as HMGCS and belongs to the HMG-CoA synthase family. The gene encodes a 520 amino acid protein which catalyzes the condensation of acetyl-CoA and acetoacetyl-CoA into 3-hydroxy-3-methylglutaryl CoA and participates in the mevalonate pathway but its end product is cholesterol(PMID:20346956). The human and hamster enzymes are distinguished by differential inhibition by magnesium ion(PMID:2870496).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Jiang-Yun Peng	36307901	Adv Sci (Weinh)	WB
Feng Zhao	34803510	Int J Biol Sci	WB
Catherine I Real	24513310	Antiviral Res	WB

# Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




WB result of HMGCS1 antibody (17643-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HMGCS1 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 17643-1-AP (HMGCS1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 17643-1-AP (HMGCS1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-HMGCS1 (IP:17643-1-AP, 3ug; Detection:17643-1-AP 1:1000) with mouse liver tissue lysate 4000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using HMGCS1 antibody (17643-1-AP) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).