For Research Use Only

TNFSF8 Polyclonal antibody

Catalog Number: 17852-1-AP



Basic Information

Catalog Number: 17852-1-AP

Size:

GenBank Accession Number: BC093630 GeneID (NCBI):

350 μg/ml 944

Source: UNIPROT ID: Rabbit P32971
Isotype: Full Name:

IgG tumor necrosis factor (ligand)
Immunogen Catalog Number: superfamily, member 8

AG12303 Calculated MW:

234 aa, 26 kDa Observed MW: 35-40 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

IHC 1:20-1:200

Applications

Tested Applications:

IHC.ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human liver cancer tissue,

Background Information

TNFSF8, also named as CD30LG and CD153, is a member of the tumor necrosis factor (TNF) receptor superfamily, is a surface antigen used as a clinical marker for Hodgkin lymphoma and related hematologic malignancies. It induces proliferation of T-cells. CD30L enhanced the proliferation of CD3-activated T cells, but induced differential responses, including cell death, in several CD30-positive lymphoma-derived cell lines. For Glycosylation, the MW in WB detection is about 35-40 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 17852-1-AP (TNFSF8 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 17852-1-AP (TNFSF8 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).