For Research Use Only

SRP54 Polyclonal antibody

Catalog Number: 17976-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17976-1-AP BC003389
Size: GeneID (NCBI): 500 µg/ml 6729
Source: UNIPROT ID: Rabbit P61011

Isotype: Full Name:
IgG signal recognition particle 54kDa

Immunogen Catalog Number: Calculated MW: AG12166 54 kDa

12166 54 kDa
Observed MW:

Observed MV 54 kDa Purification Method:

Antigen affinity purification
Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF/ICC 1:10-1:100

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

Cited Applicatio

WB, IP

Species Specificity: human, mouse Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, NIH/3T3 cells

IP: HeLa cells.

IHC: human breast cancer tissue,

IF/ICC: HepG2 cells,

Background Information

The signal recognition particle (SRP) is a ribonucleoprotein complex that mediates the targeting of proteins to the endoplasmic reticulum (ER). The complex consists of a 7S (or 7SL) RNA and 6 different proteins, and signal recognition particle 54 (SRP54) is one of them. SRP54 binds to the signal sequence of presecretory protein as they emerge from the translating ribosomes, and then transfers them to translocating chain-associating membrane protein (TRAM). This antibody specifically recognizes the 54kd SRP54 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Piazzon Nathalie N	23221635	Nucleic Acids Res	WB,IP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 17976-1-AP (SRP54 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of HepG2 cells, using SRP54 antibody 17976-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP result of anti-SRP54 (IP:17976-1-AP, 4ug; Detection:17976-1-AP 1:500) with HeLa cells lysate 1600ug.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 17976-1-AP (SRP54 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 17976-1-AP (SRP54 Antibody) at dilution of 1:50 (under 40x lens).