For Research Use Only

IGFBP4 Polyclonal antibody Catalog Number: 18500-1-AP Featured Product 9

Featured Product 9 Publications




Basic Information	Catalog Number: 18500-1-AP	GenBank Accession Number: BC016041		Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI)	:	Recommended Dilutions:
	400 µ g/ml	3487		WB 1:500-1:3000
	Source: Rabbit	UNIPROT ID: P22692		IHC 1:50-1:500 IF/ICC 1:10-1:100
	Isotype: IgG Immunogen Catalog Number:	Full Name: insulin-like growth factor binding protein 4		
	AG13347	Calculated MW: 28 kDa Observed MW: 28-40 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, ELISA WB: C6 d Cited Applications:			
	WB, IHC, IF		IHC : human liver tissue, human colon tissue	
	Species Specificity: human, rat	IF/ICC : C6 cells,		
	Cited Species: human, rat, chicken			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	IGFBP4, also named as IBP4, is a 258 amino acid protein, which contains one IGFBP N-terminal domain and one thyroglobulin type-1 domain. IGFBP4 is the smallest member of the insulin-like growth factor binding protein family (IGFBP). It is a hepatic protein that plays a role in modulating the activity and bioavailability of IGF-I. Th expression of IGFBP4 was found to increase under conditions of hypoxia. IGFBP4 is highly expressed in adipocyt and osteoblasts and is inhibitory of IGFs in vitro. (PMID: 28938423, PMID: 34646401). IGFBP4 is a secreted protein and calculated molecular weight is 28 kDa.			
Background Information	thyroglobulin type-1 domain. IGF family (IGFBP). It is a hepatic prot expression of IGFBP4 was found to and osteoblasts and is inhibitory of	BP4 is the smallest ein that plays a role o increase under cor of IGFs in vitro. (PM	member of the insuli e in modulating the a nditions of hypoxia. I	n-like growth factor binding protein ttivity and bioavailability of IGF-I. T GFBP4 is highly expressed in adipocy
	thyroglobulin type-1 domain. IGF family (IGFBP). It is a hepatic prot expression of IGFBP4 was found to and osteoblasts and is inhibitory of and calculated molecular weight	BP4 is the smallest ein that plays a role o increase under co of IGFs in vitro. (PM is 28 kDa.	member of the insuli e in modulating the a nditions of hypoxia. I	n-like growth factor binding protein ttivity and bioavailability of IGF-I. T JFBP4 is highly expressed in adipocy 4646401). IGFBP4 is a secreted prote
	thyroglobulin type-1 domain. IGF family (IGFBP). It is a hepatic prot expression of IGFBP4 was found to and osteoblasts and is inhibitory of and calculated molecular weight Author	BP4 is the smallest ein that plays a role o increase under cor of IGFs in vitro. (PM	member of the insuli e in modulating the a nditions of hypoxia. IO ID: 28938423, PMID: 3	n-like growth factor binding protein ttivity and bioavailability of IGF-I. T GFBP4 is highly expressed in adipocy
	thyroglobulin type-1 domain. IGF family (IGFBP). It is a hepatic prot expression of IGFBP4 was found to and osteoblasts and is inhibitory of and calculated molecular weight Author Wenwen Guo	BP4 is the smallest ein that plays a role o increase under co of IGFs in vitro. (PM is 28 kDa. Pubmed ID	member of the insuli e in modulating the a nditions of hypoxia. I ID: 28938423, PMID: 3 Journal	n-like growth factor binding protein ttivity and bioavailability of IGF-I. T JFBP4 is highly expressed in adipocy 4646401). IGFBP4 is a secreted prote Application
Background Information Notable Publications	thyroglobulin type-1 domain. IGF family (IGFBP). It is a hepatic prot expression of IGFBP4 was found to and osteoblasts and is inhibitory of and calculated molecular weight Author Wenwen Guo	BP4 is the smallest ein that plays a role o increase under co of IGFs in vitro. (PM is 28 kDa. Pubmed ID 34888242	member of the insuli e in modulating the a nditions of hypoxia. IO ID: 28938423, PMID: 3 Journal Front Oncol	n-like growth factor binding protein ctivity and bioavailability of IGF-I. T GFBP4 is highly expressed in adipocy 4646401). IGFBP4 is a secreted prote Application IHC

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




C6 cells were subjected to SDS PAGE followed by western blot with 18500-1-AP (IGFBP4 Antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 18500-1-AP (IGFBP4 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 18500-1-AP (IGFBP4 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of C6 cells using 18500-1-AP (IGFBP4 antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).