

For Research Use Only

TXK Polyclonal antibody

Catalog Number: 18981-1-AP **1 Publications**



Basic Information

Catalog Number: 18981-1-AP	GenBank Accession Number: NM_003328	Purification Method: Antigen affinity purification
Size: 550 µg/ml	GeneID (NCBI): 7294	Recommended Dilutions: IHC 1:20-1:200
Source: Rabbit	UNIPROT ID: P42681	IF/ICC 1:10-1:100
Isotype: IgG	Full Name: TXK tyrosine kinase	
	Calculated MW: 61 kDa	

Applications

Tested Applications: IHC, IF/ICC, ELISA	Positive Controls: IHC : human lymphoma tissue, IF/ICC : Hela cells, HepG2 cells
Cited Applications: IHC	
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

TXK, also named as PTK4, is an enzyme that transfer a phosphate group from a phosphate donor, generally the phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation.

Notable Publications

Author	Pubmed ID	Journal	Application
Yunshu Zhu	37760841	Biomedicines	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human lymphoma using 18981-1-AP (TXK antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human lymphoma using 18981-1-AP (TXK antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of Hela cells, using TXK antibody 18981-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).