

For Research Use Only

# IKZF3 Polyclonal antibody

Catalog Number: 19055-1-AP **4 Publications**



## Basic Information

<b>Catalog Number:</b> 19055-1-AP	<b>GenBank Accession Number:</b> BC032707	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 300 µg/ml	<b>GeneID (NCBI):</b> 22806	<b>Recommended Dilutions:</b> WB 1:500-1:2000 IHC 1:20-1:200 IF/ICC 1:10-1:100
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q9UKT9	
<b>Isotype:</b> IgG	<b>Full Name:</b> IKAROS family zinc finger 3 (Aiolos)	
<b>Immunogen Catalog Number:</b> AG5537	<b>Calculated MW:</b> 509 aa, 58 kDa	
	<b>Observed MW:</b> 58-70 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IF/ICC, ELISA	<b>Positive Controls:</b> WB : HepG2 cells, Jurkat cells
<b>Cited Applications:</b> WB, IHC, IF	<b>IHC :</b> human spleen tissue,
<b>Species Specificity:</b> human	<b>IF/ICC :</b> HepG2 cells,
<b>Cited Species:</b> human, mouse	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

IKZF3, also named as Zinc finger protein Aiolos or ZNFN1A3, is a 509 amino acid protein, which contains 6 C2H2-type zinc fingers and belongs to the Ikaros C2H2-type zinc-finger protein family. IKZF3 localizes in the nucleus and cytoplasm. IKZF3 can form homodimer, and heterodimer with other IKAROS family members. IKZF3 is expressed most strongly in peripheral blood leukocytes, the spleen, and the thymus. IKZF3 as a transcription factor plays an important role in the regulation of lymphocyte differentiation. The calculated molecular weight of IKZF3 is 58 kDa, but the modified IKZF3 protein is about 65-70 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Daisuke Harama	34117218	Cell Death Discov	WB
Jung-Jyh Hung	30816208	Sci Rep	WB,IHC
Xiaoqi Shi	38758229	J Clin Immunol	IF


## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 19055-1-AP (IKZF3 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human spleen using 19055-1-AP (IKZF3 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human spleen using 19055-1-AP (IKZF3 antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of HepG2 cells, using IKZF3 antibody 19055-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).