

For Research Use Only

TINAG-Specific Polyclonal antibody



Catalog Number: 19747-1-AP

1 Publications

Basic Information

Catalog Number:

19747-1-AP

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_014464

GeneID (NCBI):

27283

UNIPROT ID:

Q9UJW2

Full Name:

tubulointerstitial nephritis antigen

Calculated MW:

55 kDa

Observed MW:

55 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:20-1:200

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

IHC

Species Specificity:

human, mouse

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse small intestine tissue, human kidney tissue, human placenta tissue

IHC : human colon tissue,

Background Information

TINAG belongs to the peptidase C1 family. TINAG mediates adhesion of proximal tubule epithelial cells via integrins alpha3-beta1 and alphaV-beta3. It is a non catalytic peptidase C1 family protein. Antibodies against TINAG are found in sera of patients with tubulointerstitial nephritis, a rare autoimmune disorder that causes acute and chronic renal injury. It has 2 isoforms produced by alternative splicing. The antibody is specific to TINAG.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuan Zhou	38617519	Transl Cancer Res	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



mouse small intestine tissue were subjected to SDS PAGE followed by western blot with 19747-1-AP (TINAG-Specific antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon using 19747-1-AP (TINAG-Specific antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human colon using 19747-1-AP (TINAG-Specific antibody) at dilution of 1:100 (under 40x lens).