For Research Use Only

BTF3L3-Specific Polyclonal antibody

Catalog Number:19753-1-AP





Basic Information	Catalog Number: 19753-1-AP	GenBank Accession Numbe Q13892	er: Purification Method: Antigen affinity purification
	Size: 260 µg/ml	GenelD (NCBI): 132556	Recommended Dilutions: WB 1:500-1:1000
	Source: Rabbit	Full Name:IHC 1:20-1:200basic transcription factor 3, like 3	
	Isotype: IgG	Calculated MW: 23 kDa	
		Observed MW: 25-30 kDa	
Applications	Tested Applications: IHC, WB,ELISA	Positive Controls:	
	Species Specificity:		: human brain tissue, : human cervical cancer tissue, human placenta
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	BTF3L3 belongs to the NAC-beta family. It is a general transcription factor. It is required for the initiation of transcription. The antibody has no cross reaction to BTF3.		
Storage	Storage: Store at -20°C. Stable for one y Storage Buffer: PBS with 0.02% sodium azide Aliquoting is unnecessary for	and 50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**





human brain tissue were subjected to SDS PAGE followed by western blot with 19753-1-AP (BTF3L3-Specific antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human cervical cancer using 19753-1-AP (BTF3L3-Specific antibody) at dilution of 1:50 (under 40x lens). Immunohistochemical analysis of paraffinembedded human cervical cancer using 19753-1-AP (BTF3L3-Specific antibody) at dilution of 1:50 (under 10x lens).