

For Research Use Only

TM7SF3 Polyclonal antibody

Catalog Number: 19826-1-AP



Basic Information

Catalog Number: 19826-1-AP	GenBank Accession Number: BC005176	Purification Method: Antigen affinity purification
Size: 600 µg/ml	GeneID (NCBI): 51768	Recommended Dilutions: WB 1:500-1:3000 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9NS93	
Isotype: IgG	Full Name: transmembrane 7 superfamily member 3	
Immunogen Catalog Number: AG13919	Calculated MW: 570 aa, 64 kDa	
	Observed MW: 64-80 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls:
Species Specificity: human, mouse	WB: mouse kidney tissue, HepG2 cells, mouse pancreas tissue, mouse testis tissue, U2OS cells
	IF/ICC: U-251 cells,

Background Information

The seven-transmembrane superfamily member 3 protein (TM7SF3) is a pro-survival factor induced by p53 that regulates protein homeostasis and attenuates cellular stress. TM7SF3 inhibits caspase 3/7 while siRNA-mediated silencing of TM7SF3 accelerates ER stress and the unfolded protein responses (UPR). This involves inhibitory phosphorylation of eIF2 α activity and increased expression of ATF3, ATF4, and C/EBP, followed by induction of apoptosis. TM7SF3 maintains cellular reducing power within physiological levels and reduces the content of proapoptotic proteins such as FAS, FADD, and caspase-8. (PMID: 36304109)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 19826-1-AP (TM7SF3 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 19826-1-AP (TM7SF3 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed U-251 cells using TM7SF3 antibody (19826-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).