For Research Use Only

## MPRIP Polyclonal antibody

Catalog Number:20040-1-AP <u>3 Publications</u>

Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 20040-1-AP	GenBank Ac NM_015134	ccession Number:	Purification Method: Antigen affinity purification	
	Size:	e: GenelD (NCBI):		Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	400 µ g/ml 23164				
	Source: Rabbit	UNIPROT ID Q6WCQ1	r.	protein lysate IHC 1:50-1:500 IF 1:10-1:100	
	lsotype: IgG	Full Name: myosin pho	sphatase Rho interacting		
		protein	protein		
	Calculated MW: 117 kDa Observed MW: 116-140 kDa		MW:		
Applications	Tested Applications:		Positive Controls:		
	IF/ICC, IHC, IP, WB, ELISA WB : 5 Cited Applications:		WB: SH-SY5Y	I-SY5Y cells, A2780 cells, SKOV-3 cells	
	WB, IF	P SKUV-3 C			
	Species Specificity: IHC : mouse   human, mouse, rat IF : Hela cel		wary tissue, human ovary tissue		
	Cited Species: human				
	Note-IHC: suggested ant TE buffer pH 9.0; (*) Alte retrieval may be perforn buffer pH 6.0	ernatively, antige			
Background Information	MPRIP, also named as KIAA0864, MRIP, RHOIP3 and p116Rip, is required for the regulation of the actin cytoskeleto by RhoA and ROCK1. MRPIP targets myosin phosphatase to the actin cytoskeleton. Depletion leads to an increase number of stress fibers in smooth muscle cells through stabilization of actin fibers by phosphorylated myosin. Overexpression of MRIP as well as its F-actin-binding region leads to disassembly of stress fibers in neuronal cell The antibody recognizes the C-term of isoform2 and isoform3 of MPRIP.				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Нао Во	30588252	J Cancer	WB	
	Shan Wang	32530539	Glia	WB,IF	
	Maya David	35961752	RNA	WB	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 20040-1-AP (MPRIP antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human ovary using 20040-1-AP (MPRIP antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human ovary using 20040-1-AP (MPRIP antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of Hela cells, using MPRIP antibody 20040-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP result of anti-MPRIP (IP:20040-1-AP, 4ug; Detection:20040-1-AP 1:500) with SKOV-3 cells lysate 2000ug.



Immunohistochemical analysis of paraffinembedded mouse ovary tissue slide using 20040-1-AP (MPRIP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).