

For Research Use Only

p300 Polyclonal antibody

Catalog Number: 20695-1-AP **1 Publications**



Basic Information

Catalog Number: 20695-1-AP	GenBank Accession Number: NM_001429	Purification Method: Antigen affinity purification
Size: 500 µg/ml	GeneID (NCBI): 2033	Recommended Dilutions: WB 1:500-1:1000 IHC 1:250-1:1000
Source: Rabbit	UNIPROT ID: Q09472	
Isotype: IgG	Full Name: E1A binding protein p300	
	Calculated MW: 264 kDa	
	Observed MW: 300 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HeLa cells, MCF-7 cells, HEK-293T cells, Jurkat cells, PC-3 cells IHC : rat lung tissue, mouse lung tissue
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

p300 is a 300 kDa global transcriptional coactivator with histone acetyltransferase (HAT) activity. p300 binds to chromosomal histones and modulates the chromatin conformation by acetylation of its histones, leading to activation of various gene transcriptions. p300 plays a critical role in DNA repair synthesis after DNA damage induced by ultraviolet or radiation. p300 is overexpressed in various types of cancer, which is associated with worse patient outcomes (PMID: 27322077).

Notable Publications

Author	Pubmed ID	Journal	Application
Yinghong Zhu	38113887	Cell Metab	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 20695-1-AP (p300 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 20695-1-AP (p300 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 20695-1-AP (p300 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat lung tissue slide using 20695-1-AP (p300 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).