For Research Use Only

GABPA Polyclonal antibody Catalog Number: 21542-1-AP Featured Product

Featured Product 12 Publications




	Catalog Number: 21542-1-AP	GenBank Accession Number: BC035031	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	500 µg/ml	2551	WB 1:2000-1:16000	
	Source:	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Rabbit	Q06546	IHC 1:50-1:500	
	Isotype: IgG	Full Name: GA binding protein transcription	IF/ICC 1:200-1:800	
	Immunogen Catalog Number:	factor, alpha subunit 60kDa		
	AG16191	Calculated MW: 454 aa, 51 kDa		
		Observed MW: 56-60 kDa		
Applications	Tested Applications:	Positive 0	Positive Controls:	
	WB, IHC, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF	MCF-7 ce	WB : HEK-293 cells, A431 cells, mouse liver tissue, MCF-7 cells, A549 cells, rat brain tissue, K-562 cells, NIH/3T3 cells, mouse brain tissue	
	Species Specificity:	IP : HeLa o	IP : HeLa cells,	
	human, mouse, rat	IHC : hum	an cervical cancer tissue, human breast	
	Cited Species:	cancer tis	sue	
	human, mouse IF/ICC : HEK-293 cells,			
	NOTE-INC: SUGGESTEG ANTIGE	n retneval with		
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0	atively, antigen		
Background Information	TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GA-binding protein alpha chain ((factor E4TF1-60) is one of three G subunit.GABPA is a member of Ets Since this subunit shares identity in activation of cytochrome oxida shares identity with a subunit cor adenovirus E4 gene. Because of it	atively, antigen d with citrate GABP alpha subunit, GABPA, nuclear re A-binding protein transcription factor s family, binds to the Yap promoter an with a subunit encoding the nuclear re	subunits which functions as a DNA-binding d activates YAP transcription(23684612). espiratory factor 2 gene, it is likely involve nitochondrial function. This subunit also 1, responsible for expression of the ty to form heterodimers with other	
	TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GA-binding protein alpha chain ((factor E4TF1-60) is one of three G subunit.GABPA is a member of Ets Since this subunit shares identity in activation of cytochrome oxida shares identity with a subunit cor adenovirus E4 gene. Because of it	atively, antigen d with citrate GABP alpha subunit, GABPA, nuclear re A-binding protein transcription factor s family, binds to the Yap promoter an with a subunit encoding the nuclear r use expression and nuclear control of r istituting the transcription factor E4TF is chromosomal localization and abili	espiratory factor 2 gene, it is likely involve nitochondrial function. This subunit also 1, responsible for expression of the ty to form heterodimers with other	
Background Information	TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GA-binding protein alpha chain ((factor E4TF1-60) is one of three G subunit.GABPA is a member of Et: Since this subunit shares identity in activation of cytochrome oxida shares identity with a subunit cor adenovirus E4 gene. Because of it polypeptides, this gene may play	atively, antigen d with citrate GABP alpha subunit, GABPA, nuclear re A-binding protein transcription factor s family, binds to the Yap promoter an with a subunit encoding the nuclear r use expression and nuclear control of r istituting the transcription factor E4TF is chromosomal localization and abili a role in the Down Syndrome phenoty	subunits which functions as a DNA-binding d activates YAP transcription(23684612). espiratory factor 2 gene, it is likely involve nitochondrial function. This subunit also 1, responsible for expression of the ty to form heterodimers with other ype.	
	TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 GA-binding protein alpha chain (0 factor E4TF1-60) is one of three G subunit.GABPA is a member of Et Since this subunit shares identity in activation of cytochrome oxida shares identity with a subunit cor adenovirus E4 gene. Because of it polypeptides, this gene may play Author	Atively, antigen d with citrate GABP alpha subunit, GABPA, nuclear re A-binding protein transcription factor s family, binds to the Yap promoter an with a subunit encoding the nuclear r with a subunit encoding the nuclear r siste expression and nuclear control of r isstituting the transcription factor E4TF iss chromosomal localization and abili a role in the Down Syndrome phenoty Pubmed ID Journal	subunits which functions as a DNA-binding d activates YAP transcription(23684612). espiratory factor 2 gene, it is likely involve nitochondrial function. This subunit also 1, responsible for expression of the ty to form heterodimers with other ype. Application	


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 21542-1-AP (GABPA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer using 21542-1-AP (GABPA antibody) at dilution of 1:50 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 21542-1-AP (GABPA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 21542-1-AP (GABPA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-GABPA (IP:21542-1-AP, 4ug; Detection:21542-1-AP 1:1000) with HeLa cells lysate 1200ug.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using GABPA antibody (21542-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).