

For Research Use Only

GIRK2 Polyclonal antibody

Catalog Number: 21647-1-AP 4 Publications



Basic Information

Catalog Number:

21647-1-AP

Size:

650 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG16344

GenBank Accession Number:

BC101547

GeneID (NCBI):

3763

UNIPROT ID:

P48051

Full Name:

potassium inwardly-rectifying channel, subfamily J, member 6

Calculated MW:

423 aa, 48 kDa

Observed MW:

45-48 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IHC: mouse brain tissue,

Background Information

GIRK2 (also known as Kir3.2), encoded by KCNJ6 gene, is a member of the G protein-coupled inwardly-rectifying potassium channel (GIRK, Kir3) family of inward rectifier potassium channels. GIRK channels are activated following stimulation of G protein-coupled receptors (PMID: 26422984). They play important roles in regulating cellular excitabilities in the heart and brain (PMID: 31043612). Mutations in KCNJ6 gene are associated with Keppen-Lubinsky Syndrome (PMID: 25620207).

Notable Publications

Author	Pubmed ID	Journal	Application
Jiaxun Nie	32683743	FASEB J	WB
Colleen S Stein	35280925	Mol Ther Nucleic Acids	WB
Peiwu Ye	37739206	Prog Neurobiol	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21647-1-AP (GIRK2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21647-1-AP (GIRK2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 21647-1-AP (GIRK2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).