For Research Use Only

## YY1 Polyclonal antibody Catalog Number:22156-1-AP Featured Produce

Featured Product






Basic Information	Catalog Number: 22156-1-AP	GenBank Accession Number: BC 037308	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	500 µg/ml	7528	WB 1:500-1:2000	
	Source: Rabbit	UNIPROT ID: P25490	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	lsotype:	Full Name:	IHC 1:50-1:500	
	lgG	YY1 transcription factor	IF/ICC 1:50-1:500	
	Immunogen Catalog Number: AG17792	Calculated MW: 414 aa, 45 kDa		
		Observed MW: 65-70 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA Cited Applications:	WB : Ju	ırkat cells, HeLa cells, A549 cells, HEK-293 cel	
	WB, IHC, IF, IP, CoIP, chIP	IP : He	cells, Ian breast cancer tissue, IepG2 cells,	
	Species Specificity: human, mouse			
	Cited Species:			
	human, mouse, rat, pig, bovine			
	Note-IHC: suggested antigen retrieval with <b>TE buffer pH 9.0; (*)</b> Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Informatio	<ul> <li>buffer pH 6.0</li> <li>YY1, also named as DELTA, INO80 transcription factor family. YY1 is on a large number of cellular and direct histone deacetylases and h thus implicating histone modifica of 414 amino acids with a predict protein, probably due to the struct</li> </ul>	DS and NF-E1, contains four C2H2-ty a multifunctional transcription fac viral genes by binding to sites ove istone acetyltransferases to a pron ation in the YY1. The open reading 1 ed molecular weight of 44 kDa. Ho ture of the protein. It is a ubiquitou esis, differentiation, replication an	rlapping the transcription start site. YY1 may noter in order to activate or repress the promo frame of the human YY1 cDNA encodes a prote	
	<ul> <li>buffer pH 6.0</li> <li>YY1, also named as DELTA, INO80 transcription factor family. YY1 is on a large number of cellular and direct histone deacetylases and h thus implicating histone modifica of 414 amino acids with a predict protein, probably due to the struct fundamental roles in embryogene antibody raised against full lengt</li> </ul>	DS and NF-E1, contains four C2H2-ty a multifunctional transcription fac viral genes by binding to sites ove istone acetyltransferases to a pron ation in the YY1. The open reading 1 ed molecular weight of 44 kDa. Ho ture of the protein. It is a ubiquitou esis, differentiation, replication an th YY1 of human origin.	tor that exhibits positive and negative control or apping the transcription start site. YY1 may noter in order to activate or repress the promot frame of the human YY1 cDNA encodes a prote wever, YY1 migrates on SDS gels as a 65-68 kI sly expressed transcription factor with d proliferation. This is a rabbit polyclonal	
	<ul> <li>buffer pH 6.0</li> <li>YY1, also named as DELTA, INO80 transcription factor family. YY1 is on a large number of cellular and direct histone deacetylases and h thus implicating histone modifica of 414 amino acids with a predict protein, probably due to the struct fundamental roles in embryogene antibody raised against full lengt</li> <li>Author</li> </ul>	DS and NF-E1, contains four C2H2-ty a multifunctional transcription fac viral genes by binding to sites ove istone acetyltransferases to a pron ation in the YY1. The open reading f ed molecular weight of 44 kDa. Ho ture of the protein. It is a ubiquitou esis, differentiation, replication an th YY1 of human origin.	tor that exhibits positive and negative contro or apping the transcription start site. YY1 may noter in order to activate or repress the promo frame of the human YY1 CDNA encodes a prote wever, YY1 migrates on SDS gels as a 65-68 kd sly expressed transcription factor with d proliferation. This is a rabbit polyclonal Application	
Background Informatio	<ul> <li>buffer pH 6.0</li> <li>YY1, also named as DELTA, INO80 transcription factor family. YY1 is on a large number of cellular and direct histone deacetylases and h thus implicating histone modifica of 414 amino acids with a predict protein, probably due to the struct fundamental roles in embryogen antibody raised against full lengt</li> <li>Author Trevor Teafatiller</li> </ul>	DS and NF-E1, contains four C2H2-ty a multifunctional transcription fac viral genes by binding to sites ove istone acetyltransferases to a pron ation in the YY1. The open reading f ed molecular weight of 44 kDa. Ho ture of the protein. It is a ubiquitou esis, differentiation, replication an th YY1 of human origin. Pubmed ID Journal 36096242 Life Sci	tor that exhibits positive and negative contro relapping the transcription start site. YY1 may noter in order to activate or repress the promo frame of the human YY1 cDNA encodes a prote wever, YY1 migrates on SDS gels as a 65-68 kl sly expressed transcription factor with d proliferation. This is a rabbit polyclonal Application WB	
	<ul> <li>buffer pH 6.0</li> <li>YY1, also named as DELTA, INO80 transcription factor family. YY1 is on a large number of cellular and direct histone deacetylases and h thus implicating histone modifica of 414 amino acids with a predict protein, probably due to the struct fundamental roles in embryogene antibody raised against full lengt</li> <li>Author Trevor Teafatiller</li> <li>Qing-Dong Wang</li> </ul>	DS and NF-E1, contains four C2H2-ty a multifunctional transcription fac viral genes by binding to sites ove istone acetyltransferases to a pron ation in the YY1. The open reading t ed molecular weight of 44 kDa. Ho ture of the protein. It is a ubiquitou esis, differentiation, replication an th YY1 of human origin. Pubmed ID Journal 36096242 Life Sci 36285444 Pathol Int	tor that exhibits positive and negative control or lapping the transcription start site. YY1 may noter in order to activate or repress the promot frame of the human YY1 cDNA encodes a prote wever, YY1 migrates on SDS gels as a 65-68 kG sly expressed transcription factor with d proliferation. This is a rabbit polyclonal Application WB WB,CoIP	
	<ul> <li>buffer pH 6.0</li> <li>YY1, also named as DELTA, INO80 transcription factor family. YY1 is on a large number of cellular and direct histone deacetylases and h thus implicating histone modifica of 414 amino acids with a predict protein, probably due to the struct fundamental roles in embryogen antibody raised against full lengt</li> <li>Author Trevor Teafatiller</li> </ul>	DS and NF-E1, contains four C2H2-ty a multifunctional transcription fac viral genes by binding to sites ove istone acetyltransferases to a pron ation in the YY1. The open reading f ed molecular weight of 44 kDa. Ho ture of the protein. It is a ubiquitou esis, differentiation, replication an th YY1 of human origin. Pubmed ID Journal 36096242 Life Sci	tor that exhibits positive and negative contro relapping the transcription start site. YY1 may noter in order to activate or repress the promo frame of the human YY1 cDNA encodes a prote wever, YY1 migrates on SDS gels as a 65-68 kl sly expressed transcription factor with d proliferation. This is a rabbit polyclonal Application WB	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 22156-1-AP (YY1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Jurkat cells were subjected to SDS PAGE followed by western blot with 22156-1-AP (YY1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



A549 cells (shcontrol and shRNA of YY1) were subjected to SDS PAGE followed by western blot with 22156-1-AP (YY1 antibody) at dilution of 1:1000.



IP result of anti-YY1 (IP:22156-1-AP, 5ug; Detection:22156-1-AP 1:700) with HeLa cells lysate 880ug.



Various lysates were subjected to SDS PAGE followed by western blot with 22156-1-AP (YY1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using YY1 antibody (22156-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), Alpha Tubulin antibody (66031-1-Ig, Clone: 1E4C11, red).