## For Research Use Only

## CNKSR2 Polyclonal antibody

Catalog Number: 24054-1-AP



**Basic Information** 

Catalog Number:

24054-1-AP Size: 350 μg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG21183

Observed MW: 110 kDa, 130 kDa

Species Specificity:

**Purification Method:** Antigen Affinity purified Recommended Dilutions: WB 1:500-1:2000

**Tested Applications:** 

WB, ELISA

human, mouse, rat

Positive Controls:

WB: mouse brain tissue, rat brain tissue

## **Background Information**

**Applications** 

The Connector Enhancer of Kinase Suppressor of Ras-2 (CNKSR2), also known as CNK2 or MAGUIN, KIAA0902  $and KSR2, is a scaffolding \ molecule \ that \ contains \ functional \ protein \ binding \ domains \ (PMID: 35053419). \ CNKSR2$ interacts with different molecules, including RAF1, ARHGAP39, and CYTH2, and regulates the Mitogen-Activated Protein Kinase (MAPK) cascade and small GTPase signaling (PMID: 35053419). The study identify a molecular complex comprising CNKSR2 and the guanine nucleotide exchange factor (GEF) for ARF small GTPases, CYTH2, that is necessary for the proper development of granule neurons in the mouse hippocampus (PMID: 34800437). CNKSR2 plays a key role in regulating cell proliferation, and differentiation through multiple receptor tyrosine kinase pathways (PMID: 35053419). The molecular weight of CNKSR2 is mainly 117 kDa (PMID: 29534682). It has also been reported that CNKSR2 immunoblotted signal of about 110 kDa was detected, while a 130 kDa signal was detected (PMID: 34800437).

GenBank Accession Number:

connector enhancer of kinase suppressor of Ras 2

BC126121

22866

Q8WXI2 Full Name:

GeneID (NCBI):

**UNIPROT ID:** 

Calculated MW: 1034 aa. 118 kDa

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24054-1-AP (CNKSR2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.