For Research Use Only

SORBS2 Polyclonal antibody

Catalog Number:24643-1-AP

Featured Product

9 Publications




Basic Information	Catalog Number: 24643-1-AP	GenBank Accession Number: BC011883		Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):		Recommended Dilutions:	
	800 µ g/ml	8470		WB 1:1000-1:6000	
	Source:	UNIPROT ID:		IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Rabbit 094875				
	Isotype:	Full Name:		IHC 1:250-1:1000	
	IgG sorbin and SH3 domain containin		domain containing 2	g 2	
	Immunogen Catalog Number: AG20041		Calculated MW: 1100 aa, 124 kDa Observed MW:		
		Observed MW:			
		70-75 kDa	70-75 kDa		
Applications	Tested Applications:	Positive Controls:			
	WB, IP, IHC, ELISA		WB: A2780 ce	WB : A2780 cells, MDCK cells, U-251 cells, mouse pancreas tissue, rat pancreas tissue	
	•		pancreas tissu		
			IP : mouse hea	ouse heart tissue,	
	Species Specificity: human, mouse, canine		IHC : mouse h	: mouse heart tissue, human pancreas cancer	
	Cited Species:	tissue			
	human, mouse				
	with TE buffer pH 9.0; (*	*) Alternatively			
	Note-IHC: suggested ar with TE buffer pH 9.0; (* antigen retrieval may b with citrate buffer pH 6.	e performed	, ,		
Background Information	antigen retrieval may be with citrate buffer pH 6. SORBS2, also known as ARGBP2, h domain that interacts with lipid ra abundant in the heart. In epithelia SORBS2 is located in the Z-disks of adapter protein to assemble signa	e performed of has three C-terminal aft proteins. SORBS2 al cells SORBS2 is loo of sarcomeres. The su aling complexes in s	SH3 domains and an is widely expressed i cated in stress fibers a ubcellular localizatior tress fibers, and may	and the nucleus; In cardiac muscle ce n suggests that SORBS2 functions as a influence the contractile or elastic	
	antigen retrieval may be with citrate buffer pH 6. SORBS2, also known as ARGBP2, h domain that interacts with lipid ra abundant in the heart. In epithelia SORBS2 is located in the Z-disks of adapter protein to assemble signa properties of cardiac sarcomeres(28961272).	e performed of has three C-terminal aft proteins. SORBS2 al cells SORBS2 is low of sarcomeres. The su aling complexes in s PMID: 9211900). Alte	SH3 domains and an is widely expressed i cated in stress fibers a ubcellular localizatior tress fibers, and may	n human tissues and extremely and the nucleus; In cardiac muscle ce n suggests that SORBS2 functions as a	
	antigen retrieval may be with citrate buffer pH 6. SORBS2, also known as ARGBP2, f domain that interacts with lipid r abundant in the heart. In epithelia SORBS2 is located in the Z-disks of adapter protein to assemble signa properties of cardiac sarcomeres(28961272).	e performed O has three C-terminal aft proteins. SORBS2 al cells SORBS2 is lo of sarcomeres. The su aling complexes in s PMID: 9211900). Alte Pubmed ID	SH3 domains and an is widely expressed i cated in stress fibers a ubcellular localization tress fibers, and may ernate splicing results	In human tissues and extremely and the nucleus; In cardiac muscle ce n suggests that SORBS2 functions as a influence the contractile or elastic in multiple transcript variants(PMIC	
	antigen retrieval may be with citrate buffer pH 6. SORB52, also known as ARGBP2, H domain that interacts with lipid rr abundant in the heart. In epithelia SORB52 is located in the Z-disks of adapter protein to assemble signs properties of cardiac sarcomeres(28961272). Author Praju Vikas Anekal	has three C-terminal aft proteins. SORBS2 al cells SORBS2 is loo of sarcomeres. The su aling complexes in s PMID: 9211900). Alter Pubmed ID 25429109	SH3 domains and an is widely expressed i cated in stress fibers a ubcellular localization tress fibers, and may ernate splicing results Journal	In human tissues and extremely and the nucleus; In cardiac muscle cen suggests that SORBS2 functions as a influence the contractile or elastic in multiple transcript variants(PMIC Application	
Background Information	Antigen retrieval may be with citrate buffer pH 6. SORBS2, also known as ARGBP2, f domain that interacts with lipid ra abundant in the heart. In epithelia SORBS2 is located in the Z-disks of adapter protein to assemble signa properties of cardiac sarcomeres(28961272). Author Praju Vikas Anekal Ming-Ming Liu	e performed o has three C-terminal aft proteins. SORBS2 al cells SORBS2 is low of sarcomeres. The su- aling complexes in s PMID: 9211900). Alter Pubmed ID 25429109 35590369	SH3 domains and an is widely expressed i cated in stress fibers a ubcellular localizatior tress fibers, and may ernate splicing results Journal J Biol Chem	In human tissues and extremely and the nucleus; In cardiac muscle ce in suggests that SORBS2 functions as a influence the contractile or elastic in multiple transcript variants(PMID Application WB, IP WB	
	Antigen retrieval may be with citrate buffer pH 6. SORBS2, also known as ARGBP2, f domain that interacts with lipid ra abundant in the heart. In epithelia SORBS2 is located in the Z-disks of adapter protein to assemble signa properties of cardiac sarcomeres(28961272). Author Praju Vikas Anekal Ming-Ming Liu	e performed of has three C-terminal aft proteins. SORBS2 al cells SORBS2 is low of sarcomeres. The su aling complexes in s PMID: 9211900). Alter Pubmed ID 25429109 35590369 25818930 r after shipment.	SH3 domains and an is widely expressed i cated in stress fibers a bocellular localization tress fibers, and may ernate splicing results Journal J Biol Chem J Transl Med J Cardiovasc Pharma	In human tissues and extremely and the nucleus; In cardiac muscle ce in suggests that SORBS2 functions as a influence the contractile or elastic in multiple transcript variants(PMID Application WB, IP WB	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 24643-1-AP (SORBS2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-SORBS2 (IP:24643-1-AP, 4ug; Detection:24643-1-AP 1:500) with mouse heart tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 24643-1-AP (SORBS2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 24643-1-AP (SORBS2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).