For Research Use Only

TNN Polyclonal antibody Catalog Number:24839-1-AP




Basic Information	Catalog Number: 24839-1-AP	GenBank Accession Number: BC 136619	Purification Method: Antigen affinity purification
	Size: 550 ug/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG18127	GenelD (NCBI): 63923	Recommended Dilutions: WB 1:500-1:2000
		UNIPROT ID: Q9UQP3	
		Full Name: tenascin N	
		Calculated MW: 1299 aa, 144 kDa	
		Observed MW: 144 kDa, 173 kDa	
Applications	Tested Applications:	Positive C	Controls:
	WB, ELISA Species Specificity: human	WB : mous	se brain tissue, rat brain tissue
Background Information	TNN (Tenascin-W), also known as TNW. It is expected to be located in the extracellular space. The protein is not detected in normal adult mammary tissues or brain but expressed in most breast tumors and brain tumors, such as glioblastomas, astrocytomas and oligodendrogliomas, tested (PMID:17909022). And in brain tumors, detected around the endothelial cell layer of the clood vessels (PMID: 19884327). TNN is the fourth and final member of the extracellular matrix protein family of tenascins. In vitro, the presence of tenascin-W increased the proportion of elongated human umbilical vein endothelial cells (HUVECs) and augmented the mean speed of cell migration. Furthermore, tenascin-W irggered sprouting of HUVEC spheroids to a similar extent as the proangiogenic factor tenascin-C. Tenascin-W is a specific marker of glioma-associated blood vessels and stimulates angiogenesis in vitro(PMID: 19884327). The predicted molecular weight of the protein is 144 kDa in human and 173 kDa in mouse.		
	Storage:		

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24839-1-AP (TNN antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.