For Research Use Only

F2RL3 Polyclonal antibody

Catalog Number:25306-1-AP 3 Publications




Basic Information	Catalog Number: 25306-1-AP	GenBank Accession Number: BC074782	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	800 µg/ml	9002	WB 1:500-1:2000
	Source: Rabbit	UNIPROT ID: Q96RIO	IHC 1:20-1:200
	Isotype: Full Name: IgG coagulation factor II (thrombin)		
	Immunogen Catalog Number: AG20801	receptor-like 3 Calculated MW:	
		385 aa, 41 kDa	
		Observed MW: 41 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, ELISA	WB : mouse lung tissue, mouse pancreas tissue	
	Cited Applications: WB	IHC : mou	se lung tissue,
	Species Specificity: human, mouse		
	Cited Species: human, mouse		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
	retrieval may be performed		
Background Information	retrieval may be performed buffer pH 6.0 Coagulation factor II receptor-like known as protease-activated rece inflammation (PMID:34284820). A protection against pulmonary em	d with citrate e 3 (F2RL3) encodes a member of the p eptor 4 (PAR4), which takes partin plate An absence of PAR4 in mouse models i bolism, and a small number of misser i have been described(PMID:35012325	elet activation, intimal hyperplasia and
	retrieval may be performed buffer pH 6.0 Coagulation factor II receptor-like known as protease-activated rece inflammation (PMID:34284820). A protection against pulmonary em platelet aggregation and functior	d with citrate e 3 (F2RL3) encodes a member of the p eptor 4 (PAR4), which takes partin plate An absence of PAR4 in mouse models i bolism, and a small number of misser i have been described(PMID:35012325	results in impaired hemostasis and a nse coding variants in F2RL3 that alter). The calculated molecular weight and
	retrieval may be performed buffer pH 6.0 Coagulation factor II receptor-like known as protease-activated rece inflammation (PMID:34284820). A protection against pulmonary em platelet aggregation and functior observed molecular weight of F20 Author	d with citrate e 3 (F2RL3) encodes a member of the p ptor 4 (PAR4), which takes partin plate An absence of PAR4 in mouse models i bolism, and a small number of misser in have been described(PMID:35012325 RL3 are both 41 kDa. Pubmed ID Journal	elet activation, intimal hyperplasia and results in impaired hemostasis and a nse coding variants in F2RL3 that alter
	retrieval may be performed buffer pH 6.0 Coagulation factor II receptor-like known as protease-activated rece inflammation (PMID:34284820). / protection against pulmonary em platelet aggregation and functior observed molecular weight of F20	d with citrate e 3 (F2RL3) encodes a member of the p ptor 4 (PAR4), which takes partin plate An absence of PAR4 in mouse models a bolism, and a small number of misser a have been described(PMID:35012325 RL3 are both 41 kDa. Pubmed ID Journal	elet activation, intimal hyperplasia and results in impaired hemostasis and a rese coding variants in F2RL3 that alter j). The calculated molecular weight and Application
	retrieval may be performed buffer pH 6.0 Coagulation factor II receptor-like known as protease-activated rece inflammation (PMID:34284820). A protection against pulmonary em platelet aggregation and function observed molecular weight of F20 Author Jianing Luo	d with citrate e 3 (F2RL3) encodes a member of the p ptor 4 (PAR4), which takes partin plate An absence of PAR4 in mouse models 1 bolism, and a small number of misser have been described(PMID:35012325 RL3 are both 41 kDa. Pubmed ID Journal 33111257 Neurosci Bull	elet activation, intimal hyperplasia and results in impaired hemostasis and a nse coding variants in F2RL3 that alter)). The calculated molecular weight and Application WB WB
Background Information	retrieval may be performed buffer pH 6.0 Coagulation factor II receptor-like known as protease-activated rece inflammation (PMID:34284820). A protection against pulmonary em platelet aggregation and functior observed molecular weight of F20 Author Jianing Luo Nathalia Araujo	a with citrate a 3 (F2RL3) encodes a member of the p ptor 4 (PAR4), which takes partin plate An absence of PAR4 in mouse models 1 bolism, and a small number of misser have been described(PMID:35012325 RL3 are both 41 kDa. Pubmed ID Journal 33111257 Neurosci Bull 35425699 Front Oncol 35082966 Oxid Med Cell Lo	elet activation, intimal hyperplasia and results in impaired hemostasis and a nse coding variants in F2RL3 that alter). The calculated molecular weight and Application WB WB


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 25306-1-AP (F2RL3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



mouse lung tissue were subjected to SDS PAGE followed by western blot with 25306-1-AP (F2RL3 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using 25306-1-AP (F2RL3 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using 25306-1-AP (F2RL3 Antibody) at dilution of 1:50 (under 40x lens).