For Research Use Only

APP/Beta Amyloid Polyclonal antibody

Catalog Number:25524-1-AP

Featured Product 101 Publications


y proteintech Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 25524-1-AP	GenBank Accessi BC065529	ion Number:	Purification Method: Antigen affinity purification	
	Concentration:	GenelD (NCBI):		Recommended Dilutions:	
	550 ug/ml	351		WB: 1:500-1:2000	
	Source:	UNIPROT ID:		IHC: 1:50-1:500	
	Rabbit	P05067		IF/ICC: 1:50-1:500	
	Isotype:	Full Name:			
	IgG	amyloid beta (A4) precursor protein Observed MW: 100 kDa			
	Immunogen Catalog Number: AG22408				
Applications	Tested Applications:	Positive Controls:			
	WB, IHC, IF/ICC, ELISA		WB : SH-SY5Y cells, HAP1 cells, HeLa cells, NCI-H129		
	Cited Applications:		cells, mouse	cells, mouse brain tissue, rat brain tissue, C6 cells	
	WB, IHC, IF		IHC : human	HC : human gliomas tissue, human brain tissue	
	Species Specificity: human, mouse, rat		IF/ICC : SH-SY5Y cells,		
	Cited Species: human, mouse, rat, zebrafish				
	hadden will C.C.				
Background Informatic	cleavage precludes the formation	of A β , while the β -	and γ -cleavages g		
Background Informatic	A β derives from APP via proteoly cleavage precludes the formation amyloidogenic features. Amyloid chromosome 21q, is a cell surface to neurite growth, neuronal adhes brain, kidney, heart and spleen, b Amyloid β (A β) precursor protei isoforms. This antibody can recog protein, including APP770, APP67 several chains, this antibody coul	n of A β , while the β - d beta A4 precursor pro- e receptor and perform sion and axonogenesis ut weak in liver. Defe- in (APP) is a 100-140 k mize several isoforms 77, APP695, APP696, Al d recognize fragment use C-terminal fragme	and γ -cleavages g otein(APP), encoded is physiological fur s. APP expressed in cts in APP are the c Da transmembrand of both mature and PP733, APP751, AP s C99, Amyloid-bet	generate APP components with d by APP gene which locate on human actions on the surface of neurons relevar all fetal tissues and is pronounced in ause of Alzheimer disease type 1 (AD1). e glycoprotein that exists as several d immature amyloid beta (A4) precursor P752, and APP639. APP can be cleaved in ta protein 42, Amyloid-beta protein 40,	
Background Informatio	A β derives from APP via proteoly cleavage precludes the formation amyloidogenic features. Amyloid chromosome 21q, is a cell surface to neurite growth, neuronal adhes brain, kidney, heart and spleen, b Amyloid β (A β) precursor protei isoforms. This antibody can recog protein, including APP770, APP67 several chains, this antibody coul C83, P3(40), C80, Gamma-secreta	n of A β , while the β - d beta A4 precursor pro- e receptor and perform sion and axonogenesis ut weak in liver. Defei n (APP) is a 100-140 k mize several isoforms 77, APP695, APP696, Al d recognize fragment use C-terminal fragme 0, C31.	and γ -cleavages g otein(APP), encoded is physiological fur s. APP expressed in cts in APP are the c Da transmembrand of both mature and PP733, APP751, AP s C99, Amyloid-bet	generate APP components with d by APP gene which locate on human actions on the surface of neurons relevar a all fetal tissues and is pronounced in ause of Alzheimer disease type 1 (AD1). e glycoprotein that exists as several d immature amyloid beta (A4) precursor P752, and APP639. APP can be cleaved in	
	A β derives from APP via proteoly cleavage precludes the formation amyloidogenic features. Amyloid chromosome 21q, is a cell surface to neurite growth, neuronal adhes brain, kidney, heart and spleen, b Amyloid β (A β) precursor protei isoforms. This antibody can recog protein, including APP770, APP67 several chains, this antibody coul C83, P3(40), C80, Gamma-secreta secretase C-terminal fragment 50	h of A β , while the β - d beta A4 precursor pro- e receptor and perform sion and axonogenesis ut weak in liver. Defe- in (APP) is a 100-140 k gnize several isoforms 77, APP695, APP696, Al d recognize fragment use C-terminal fragme 0, C31.	and γ -cleavages g otein(APP), encoded is physiological fur s. APP expressed in cts in APP are the c iDa transmembrand of both mature and PP733, APP751, AP s C99, Amyloid-bet nt 59, Gamma-secr	generate APP components with d by APP gene which locate on human actions on the surface of neurons relevar all fetal tissues and is pronounced in ause of Alzheimer disease type 1 (AD1). e glycoprotein that exists as several d immature amyloid beta (A4) precursor P752, and APP639. APP can be cleaved in ta protein 42, Amyloid-beta protein 40, etase C-terminal fragment 57, Gamma-	
	A β derives from APP via proteoly cleavage precludes the formation amyloidogenic features. Amyloid chromosome 21q, is a cell surface to neurite growth, neuronal adhes brain, kidney, heart and spleen, b Amyloid β (A β) precursor protei isoforms. This antibody can recog protein, including APP770, APP67 several chains, this antibody coul C83, P3(40), C80, Gamma-secreta secretase C-terminal fragment 50 Author	n of A β , while the β - d beta A4 precursor pro- e receptor and perform sion and axonogenesi: ut weak in liver. Defe- in (APP) is a 100-140 k mize several isoforms rg, APP695, APP696, Al d recognize fragment: se C-terminal fragme 0, C31.	and γ -cleavages g otein(APP), encoded is physiological fur s. APP expressed in cts in APP are the c Da transmembrane of both mature and PP733, APP751, AP s C99, Amyloid-bet int 59, Gamma-secr	generate APP components with d by APP gene which locate on human factions on the surface of neurons relevan a lass of Alzheimer disease type 1 (AD1). e glycoprotein that exists as several d immature amyloid beta (A4) precursor P752, and APP639. APP can be cleaved in ca protein 42, Amyloid-beta protein 40, etase C-terminal fragment 57, Gamma- Application	
	A β derives from APP via protection cleavage precludes the formation amyloidogenic features. Amyloid chromosome 21q, is a cell surface to neurite growth, neuronal adhes brain, kidney, heart and spleen, b Amyloid β (A β) precursor protei isoforms. This antibody can recog protein, including APP770, APP67 several chains, this antibody could C83, P3(40), C80, Gamma-secreta secretase C-terminal fragment 50 Author Shenya Xu	n of A β , while the β - d beta A4 precursor pro- e receptor and perform sion and axonogenesis ut weak in liver. Defect in (APP) is a 100-140 k mize several isoforms 77, APP695, APP696, Al d recognize fragment ise C-terminal fragme 0, C31. Pubmed ID 33183805	and γ -cleavages g otein(APP), encoded is physiological fur s. APP expressed in cts in APP are the c iDa transmembrane of both mature and PP733, APP751, AP s C99, Amyloid-bet nt 59, Gamma-secr Journal Sci Total Environ	generate APP components with d by APP gene which locate on human actions on the surface of neurons relevar all fetal tissues and is pronounced in ause of Alzheimer disease type 1 (AD1). e glycoprotein that exists as several d immature amyloid beta (A4) precursor P752, and APP639. APP can be cleaved in ca protein 42, Amyloid-beta protein 40, etase C-terminal fragment 57, Gamma- MB, IHC IHC,IF	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 25524-1-AP (APP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 25524-1-AP (APP, C-Terminal antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 25524-1-AP (APP, C-Terminal antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using APP/Beta Amyloid antibody (25524-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).