

For Research Use Only

# cleaved-Caspase 3 Polyclonal antibody

Catalog Number: 25546-1-AP

130 Publications



## Basic Information

**Catalog Number:**

25546-1-AP

**Size:**

320 µg/ml

**Source:**

Rabbit

**Isotype:**

IgG

**GenBank Accession Number:**

NM\_032991

**GeneID (NCBI):**

836

**UNIPROT ID:**

P42574

**Full Name:**

caspase 3, apoptosis-related cysteine peptidase

**Calculated MW:**

32 kDa

**Observed MW:**

17 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF/ICC 1:20-1:200

## Applications

**Tested Applications:**

WB, IP, IF, IHC, ELISA

**Cited Applications:**

WB, IHC, IF

**Species Specificity:**

human, mouse, rat

**Cited Species:**

human, mouse, rat, hamster, sheep

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

WB: Tunicamycin treated HepG2 cells, Tunicamycin treated HeLa cells, Staurosporine treated Jurkat cells

IP: Jurkat cells,

IHC: human skeletal muscle tissue,

IF/ICC: HepG2 cells,

## Background Information

Caspase 3, also named CPP32, SCA-1, and Apopain, belongs to the peptidase C14A family. Caspase 3 is involved in the activation cascade of caspases responsible for apoptosis execution. At the onset of apoptosis, it proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Caspase 3 cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Cleaves and activates caspase-6, -7, and -9. CASP3 is involved in the cleavage of huntingtin. The antibody can recognize active-caspase 3.

## Notable Publications

Author	Pubmed ID	Journal	Application
Samana Batool	30274346	Int J Mol Sci	WB
Genquan Yue	31702026	Mol Med Rep	WB
Kate D Meyer	31548708	Nat Methods	

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Tunicamycin treated HepG2 cells were subjected to SDS PAGE followed by western blot with 25546-1-AP (Caspase 3 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




IP result of anti-cleaved-Caspase 3 (IP:25546-1-AP, 4ug; Detection:25546-1-AP 1:300) with Jurkat cells lysate 3200ug.



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 25546-1-AP (Caspase 3 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 25546-1-AP (Caspase 3 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 25546-1-AP (Caspase 3 Antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).