

For Research Use Only

UFL1 Polyclonal antibody

Catalog Number: 26087-1-AP

Featured Product

10 Publications



Basic Information

Catalog Number: 26087-1-AP	GenBank Accession Number: BC036379	Purification Method: Antigen affinity purification
Size: 700 ug/ml	GeneID (NCBI): 23376	Recommended Dilutions: WB 1:500-1:1000 IHC 1:200-1:800
Source: Rabbit	UNIPROT ID: O94874	
Isotype: IgG	Full Name: KIAA0776	
Immunogen Catalog Number: AG23404	Observed MW: 80 kDa; 100 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HEK-293T cells, IHC : mouse liver tissue,
Cited Applications: WB, IHC, IF, IP	
Species Specificity: human, mouse	
Cited Species: human, mouse, bovine	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Meiqian Kuang	31721015	Cell Stress Chaperones	WB,IHC,IF
Chengmin Li	31078114	Mol Immunol	WB
Meiqian Kuang	32655766	Oxid Med Cell Longev	WB,IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 26087-1-AP (UFL1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 26087-1-AP (UFL1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HEK-293T cells were subjected to SDS PAGE followed by western blot with 26087-1-AP (KIAA0776 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.