

For Research Use Only

NUCB2/nesfatin-1 Polyclonal antibody

Catalog Number: 26712-1-AP

3 Publications



Basic Information

Catalog Number: 26712-1-AP	GenBank Accession Number: NM_005013	Purification Method: Antigen affinity purification
Size: 700 ug/ml	GeneID (NCBI): 4925	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:50-1:500
Source: Rabbit	Full Name: nucleobindin 2	
Isotype: IgG	Calculated MW: 50 kDa	
Immunogen Catalog Number: AG25152	Observed MW: 45-50 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:
WB, IF

Species Specificity:
human, mouse, rat

Cited Species:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse brain tissue, human milk, HEK-293T cells, rat brain tissue, HepG2 cells, mouse testis tissue

IHC : human intrahepatic cholangiocarcinoma tissue,

IF/ICC : HepG2 cells,

Background Information

NucB2, an anorexigenic molecule, is expressed mainly in the hypothalamus, particularly in the supraoptic nucleus (SON) and the paraventricular nucleus (PVN). NucB2 is also expressed in the subfornical organ (SFO). Because the SON and PVN are involved in body fluid regulation, NucB2 may be involved in dehydration-induced anorexia. NUCB2 is composed of a signal peptide of 24 amino acids in the N-terminal region and a protein structure containing 396 amino acids. Nesfatin-1 is an 82-amino-acid peptide hormone cleaved from nucleobindin2 (NUCB2).

Notable Publications

Author	Pubmed ID	Journal	Application
Wang Li	29901268	Proteomics	WB
Xue-Xue Zhu	39394127	Cell Commun Signal	WB,IF
Siyi Ning	37277807	J Transl Med	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 26712-1-AP (NUCB2/nesfatin-1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 26712-1-AP (NUCB2/nesfatin-1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 26712-1-AP (NUCB2/nesfatin-1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NUCB2/nesfatin-1 antibody (26712-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



1×10^6 HepG2 cells were intracellularly stained with 0.4 ug NUCB2/nesfatin-1 Polyclonal antibody (26712-1-AP)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).