## For Research Use Only

# Bcl-XL Polyclonal antibody

Catalog Number: 26967-1-AP

66 Publications



**Basic Information** 

Catalog Number: 26967-1-AP

Size: GeneID (NCBI):

450 μ g/ml 598 Source: UNIPROT ID:

Rabbit Q07817
Isotype: Full Name:
IgG BCL2-like 1
Immunogen Catalog Number: Calculated MW:

AG25604 26 kDa

Observed MW: 30 kDa

BC019307

GenBank Accession Number:

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

Tested Applications: FC, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity: human, mouse Cited Species:

human, rat, mouse, canine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: NIH/3T3 cells, RAW 264.7 cells

IP: NIH/3T3 cells,

 $\label{lhc:human testis tissue, human lung cancer tissue} \textbf{IHC:} human testis tissue, human lung cancer tissue}$ 

# **Background Information**

BCL2L1 is a member of the BCL-2 protein family. BCL2L1 is expressed as three isoforms, Bcl-X(L),Bcl-X(s) and Bcl-X(beta), and is located at the outer mitochondrial membrane. The Bcl-XL isoform is a 233 amino acid protein, acting as an apoptotic inhibitor. Bcl-XL can forms heterodimers with BAX, BAK or BCL2, and the heterodimerization with BAX does not seem to be required for anti-apoptotic activity. The Bcl-XS isoform is a shorter variant that is 178 amino acids in length and lacks a 63 amino acid region (amino acids 126-188), acting as an apoptotic activator. This antibody can recognize Bcl-XL, Bcl-XS and Bcl-X(Beta).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Guibin Lin	31607919	Front Pharmacol	WB
Jiangjiang Fan	30246250	J Cell Physiol	WB
Xiao-Lei Xin	31542790	Med Sci Monit	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


### **Selected Validation Data**




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 26967-1-AP (Bcl-XL antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human testis tissue slide using 26967-1-AP (Bcl-XL antibody) at dilution of 1:200 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 26967-1-AP (Bcl-XL antibody) at dilution of 1:200 (under 40x lens)



IP result of anti-Bcl-XL (IP:26967-1-AP, 4ug; Detection:26967-1-AP 1:1000) with NIH/3T3 cells lysate 2000 ug.



1X10^6 Jurkat cells were intracellularly stained with 0.2 ug Anti-Human Bcl-XL (26967-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Control Antibody. Cells were fixed with 90% MeOH.