For Research Use Only

SCRIB Polyclonal antibody

Catalog Number: 27083-1-AP

1 Publications



Purification Method:

WB 1:2000-1:12000

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: 27083-1-AP

 $\begin{array}{lll} 27083\text{-}1\text{-}AP & BC036905 \\ \text{Size:} & \text{GeneID (NCBI):} \\ 1000 \ \mu \, \text{g/ml} & 23513 \\ \text{Source:} & \text{UNIPROT ID:} \\ \text{Rabbit} & Q14160 \\ \end{array}$

Isotype: Full Name:
IgG scribbled homolog (Drosophila)

Immunogen Catalog Number:Calculated MW:AG257951630 aa, 175 kDa

Observed MW: 260 kDa

GenBank Accession Number:

Applications

Tested Applications:

WB,ELISA

Cited Applications:

WB

Species Specificity: human, mouse Cited Species: mouse Positive Controls:

WB: NIH/3T3 cells,

Background Information

SCRIB encodes a large, 260-kDa cytoplasmic scaffolding protein that comprises a large leucine-rich repeat (LRR) region and 4 PDZ domains that regulate protein-protein interactions. In mammals, cell polarity is established and maintained by at least 3 protein modules (Scrib, Crumbs, and Par). The apical (Crumbs and Par) and basolateral (Scrib) modules function in a mutually antagonistic relationship to regulate various polarization processes such as apical-basal polarity, planar cell polarity, asymmetric cell division and migration. The loss of SCRIB in malignant tumors suggest that it has potential to be a tumor suppressor. Alteration of polarity through deletion, downregulation, overexpression, and mislocalization can induce structural and functional alteration of cells that might be related to tumorigenesis. Mislocalization of SCRIB is involved in epithelial-to-mesenchymal transition (EMT) by inducing loss of E-cadherin. Deregulated SCRIB expression induces tumorigenesis in breast and liver. (PMID: 28460446, PMID: 32564009)

Notable Publications

Author	Pubmed ID	Journal	Application
Zhenyong Xiao	35509839	Oxid Med Cell Longev	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 27083-1-AP (SCRIB antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.