

For Research Use Only

TAPT1 Polyclonal antibody

Catalog Number: 27657-1-AP **1 Publications**



Basic Information

Catalog Number: 27657-1-AP	GenBank Accession Number: BC066899	Purification Method: Antigen affinity purification
Size: 350 µg/ml	GeneID (NCBI): 202018	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q6NXT6	
Isotype: IgG	Full Name: transmembrane anterior posterior transformation 1	
Immunogen Catalog Number: AG26360	Calculated MW: 567 aa, 64 kDa	
	Observed MW: 65-70 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : NIH/3T3 cells, mouse testis tissue IHC : human liver cancer tissue,
Cited Applications: IHC	
Species Specificity: Human, mouse	
Cited Species: mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Chenchen Zhou	34876557	Bone Res	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 27657-1-AP (TAPT1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27657-1-AP (TAPT1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27657-1-AP (TAPT1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).