

## Basic Information

<b>Catalog Number:</b> 27789-1-AP	<b>GenBank Accession Number:</b> BC151843	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 450 µg/ml	<b>GeneID (NCBI):</b> 3371	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:500-1:2000
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P24821	
<b>Isotype:</b> IgG	<b>Full Name:</b> tenascin C	
<b>Immunogen Catalog Number:</b> AG27122	<b>Calculated MW:</b> 2201 aa, 241 kDa	

## Applications

### Tested Applications:

IHC, WB, ELISA

### Species Specificity:

Human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB : U-87 MG cells,

IHC : human liver cancer tissue, human breast cancer tissue, human lung cancer tissue

## Background Information

Tenascin-C (TNC) is a large hexameric extracellular matrix glycoprotein of the tenascin family (PMID: 21818551). It is a multimodular protein containing multiple epidermal growth factor (EGF)-like repeats and fibronectin type III (FN III) domains (PMID: 1719530). Tenascin-C is highly expressed during embryonic development, particularly in the developing central nervous system, around motile cells and at epithelial-mesenchymal interaction sites (PMID: 25738825). In adult tissues, the expression and the distribution of TNC are typically limited under normal physiological conditions. It is upregulated during injury, inflammation, regeneration, and cancer (PMID: 7694605; 25120494). Tenascin-C is a diverse protein that can produce different functions including regulating cell adhesion, migration and proliferation.

## Storage

### Storage:


Store at -20°C. Stable for one year after shipment.

### Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



U-87 MG cells were subjected to SDS PAGE followed by western blot with 27789-1-AP (TNC antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27789-1-AP (TNC/Tenascin-C antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 27789-1-AP (TNC/Tenascin-C antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).