For Research Use Only

## RNF139 Polyclonal antibody Catalog Number: 28261-1-AP




Basic Information	Catalog Number: 28261-1-AP	GenBank Accession Number: BC 064636	Purification Method: Antigen affinity purification
	Size: 700 ug/ml	GeneID (NCBI): 11236	Recommended Dilutions: WB 1:1000-1:4000
	Source: Rabbit	UNIPROT ID: Q8WU17	
	Isotype: IgG	Full Name: ring finger protein 139	
	Immunogen Catalog Number: AG27147	Calculated MW: 664 aa, 76 kDa	
		Observed MW: 70 kDa	
Applications	Tested Applications: WB, ELISA Species Specificity: human	Positive C	Controls:
			WB : HEK-293 cells, HT-29 cells, HepG2 cells, THP-1 cells, U-87 MG cells, U2OS cells
Background Information	The RNF139 gene (ring-finger protein 139), also known as TRC8 (translocation in renal carcinoma from chromosome 8), was first identified from a family with 3;8 chromosomal translocation and hereditary renal cell carcinoma and thyroid cancer (PMID:22689053). RNF139 encodes an endoplasmic reticulum (ER) protein with 10 transmembrane segments that contain a sterol-sensing domain and a RING finger motif encoding an E3 ubiquitin ligase (PMID: 19706601). Northern blot and dot blot show that RNF139 is highly expressed in the testis, placenta, and adrenal gland and is expressed at lower levels in the heart, brain, liver, skeletal muscle, and pancreas (PMID: 9689122). It is reported that TRC8 suppresses tumorigenesis by targeting heme oxygenase-1 (HO-1), an antioxidant enzyme highly expressed in various cancers, for ubiquitination and degradation (PMID:22689053). TRC8 is also involved in cholesterol and fatty acid biosynthesis that are transcriptionally regulated by the sterol response element binding proteins (SREBPs) (PMID:17016439). Acetylation, phosphorylation, and autoubiquitination are common post-translational modifications of FNF139 protein.		
	thyroid cancer (PMID:22689053). F segments that contain a sterol-ser (PMID: 19706601). Northern blot a adrenal gland and is expressed at (PMID: 9689122). It is reported tha antioxidant enzyme highly expres is also involved in cholesterol and element binding proteins (SREBPs	nsing domain and a RING finger motif nd dot blot show that RNF 139 is highly lower levels in the heart, brain, liver, t TRC8 suppresses tumorigenesis by t seed in various cancers, for ubiquitina I fatty acid biosynthesis that are trans ) (PMID:17016439). Acetylation, phos	encoding an E3 ubiquitin ligase y expressed in the testis, placenta, and skeletal muscle, and pancreas argeting heme oxygenase-1 (HO-1), an tion and degradation (PMID:22689053). TRCE criptionally regulated by the sterol response

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 28261-1-AP (RNF 139 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.