For Research Use Only

Phospho-Caspase 9 (Ser196) Polyclonal antibody



Purification Method:

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

Catalog Number: 28794-1-AP

2 Publications

Basic Information

Catalog Number: 28794-1-AP

350 μg/ml Source: Rabbit

Isotype:

Size:

GenBank Accession Number:

P55211 Full Name:

caspase 9, apoptosis-related cysteine

peptidase 46 kDa Observed MW: 36 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

Species Specificity:

Human Cited Species: human

BC002452 GeneID (NCBI):

UNIPROT ID:

Calculated MW:

Positive Controls:

WB: Calyculin A treated HEK-293T cells,

Background Information

Caspase 9 also name as MCH6, APAF3, APAF-3, ICE-LAP6 and CASPASE-9c, is a member of the cysteine-aspartic acid protease (caspase) family. It's synthesized as a 46 kDa precursor protein which can be cleaved into a 35 kDa subunit and a 11 kDa subunit. Control of all caspases is tightly regulated by a series of phosphorylation events enacted by several different kinases. Caspase-9 is the most heavily phosphorylated of all caspases, with phosphorylation of at least 11 distinct residues in all three caspase-9 domains by nine kinases. It plays a central role in the mitochondrial or intrinsic apoptotic pathway that is engaged in response to many apoptotic stimuli. Once activated, caspase-9 cleaves and activates the effector caspases 3 and 7 to bring about apoptosis. It's reported that there is an increase in caspase 9 expression and activity in the hypoxic brain. Inhibition of Caspase 9 activity would render opportunity to treat neurological diseases such as stroke, neurodegenerative diseases or brain injury caused by hypoxia. (PMID: 19788417, PMID: 10529400, PMID: 9812896, PMID: 18840507, PMID: 29066624)

Notable Publications

Author	Pubmed ID	Journal	Application
Bing-Xin Chu	34804044	Front Immunol	WB
Tianjie Pu	38335292	Sci Adv	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated HEK-293T and calyculin A treated HEK-293T cells were subjected to SDS PAGE followed by western blot with 28794-1-AP (Phospho-Caspase 9 (Ser196) antibody) at dilution of 1:2000 incubated at $4^{\circ}\mathrm{C}$ overnight. The membrane was stripped and re-blotted with GAPDH antibody as loading control.