

For Research Use Only

Phospho-STAT1 (Tyr701) Polyclonal antibody



Catalog Number: 28979-1-AP

1 Publications

Basic Information

Catalog Number:

28979-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC002704

GeneID (NCBI):

6772

UNIPROT ID:

P42224

Full Name:

signal transducer and activator of transcription 1, 91kDa

Calculated MW:

83 kDa

Observed MW:

84-91 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IF 1:200-1:800

Applications

Tested Applications:

IF/ICC, WB, ELISA

Cited Applications:

WB

Species Specificity:

Human

Cited Species:

human

Positive Controls:

WB: IFN gamma and LPS treated THP-1 cells,

IF: IFN alpha treated HeLa cells,

Background Information

STAT1 (signal transducers and activators of transcription 1) is a member of the STAT protein family. STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. The phosphorylation of STAT1 at Tyr701 promotes the dimerization of STAT1 and its translocation into the nucleus where it binds the GAS promoter to allow the transcription of inflammatory response genes (PMID: 31408005). The transcript of STAT1 undergoes alternative splicing, resulting in two isoforms: STAT1 α (91 kDa) and STAT1 β (84 kDa) (PMID: 20159032).

Notable Publications

Author	Pubmed ID	Journal	Application
Ye Xie	37980535	J Transl Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated THP-1 cells, IFN gamma and LPS treated THP-1 cells were subjected to SDS PAGE followed by western blot with 28979-1-AP (Phospho-STAT1 (Tyr701) antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed IFN alpha treated HeLa cells using Phospho-STAT1 (Tyr701) antibody (28979-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).