

For Research Use Only

# CEBP Alpha/CEBPA Polyclonal antibody

Catalog Number: 29388-1-AP

6 Publications



## Basic Information

**Catalog Number:**

29388-1-AP

**Size:**

1500 ug/ml

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG29947

**GenBank Accession Number:**

NM\_004364

**GeneID (NCBI):**

1050

**UNIPROT ID:**

P49715

**Full Name:**

CCAAT/enhancer binding protein (C/EBP), alpha

**Calculated MW:**

38 kDa

**Observed MW:**

40-45 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

## Applications

**Tested Applications:**

WB, IP, ELISA

**Cited Applications:**

WB

**Species Specificity:**

human

**Cited Species:**

human, mouse, rat

**Positive Controls:**

WB : BGC-823 cells, LNCaP cells, THP-1 cells, U-937 cells, K-562 cells, L02 cells, HL-60 cells

IP : THP-1 cells,

## Background Information

CEBPA and its isoforms play important roles in lineage determination and gene activation in a variety of cell types by activating transcription from lineage-specific promoters. CEBPA is a DNA-binding protein that recognizes two different motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers. In hematopoiesis, C/EBPα is a key factor in driving the development of myeloid cells interacting with a variety of factors, including c-Myc, PU.1, and microRNAs. It can also form heterodimers with the related proteins CEBP-beta and CEBP-gamma. The encoded protein has been shown to bind to the promoter and modulate the expression of the gene encoding leptin which plays an important role in body weight homeostasis. CEBPA can interact with CDK2 and CDK4, thereby inhibiting these kinases and causing growth arrest in cultured cells. Several pathways have been implicated as the means by which CEBPA mediates cell cycle arrest and proliferation, including p21, cyclin-dependent kinases and the E2F complex via c-Myc. The calculated molecular weight of CEBPA is 38 kDa, but modified CEBPA is about 42 kDa (PMID: 19623175).

## Notable Publications

Author	Pubmed ID	Journal	Application
Ying Li	39316261	Cell Biochem Biophys	WB
Tian Li	39351088	Front Pharmacol	WB
Asuka Takahashi	39209927	Sci Rep	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29388-1-AP (CEBP Alpha/CEBPA antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-CEBP Alpha/CEBPA (IP:29388-1-AP, 4ug; Detection:29388-1-AP 1:4000) with THP-1 cells lysate 1520 ug.