

For Research Use Only

NCOA3 Polyclonal antibody, PBS Only

Catalog Number: 29587-1-PBS



Basic Information

Catalog Number: 29587-1-PBS	GenBank Accession Number: BC119001	Purification Method: Antigen affinity purification
Size: 1 mg/ml	GeneID (NCBI): 8202	
Source: Rabbit	UNIPROT ID: Q9Y6Q9	
Isotype: IgG	Full Name: nuclear receptor coactivator 3	
Immunogen Catalog Number: AG30633	Calculated MW: 1424 aa, 155 kDa	
	Observed MW: 150-160 kDa	

Applications

Tested Applications:
WB, IHC, Indirect ELISA

Species Specificity:
human

Background Information

NCOA3, also named as AIB1, SRC3 and TRAM1, belongs to the SRC/p160 nuclear receptor coactivator family. NCOA3 is a nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. NCOA3 plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). NCOA3 also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29587-1-AP (NCOA3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 29587-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 29587-1-AP (NCOA3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 29587-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 29587-1-AP (NCOA3 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 29587-1-PBS in a different storage buffer formulation.