

For Research Use Only

cGAS Polyclonal antibody

Catalog Number: 29958-1-AP **4 Publications**



Basic Information

Catalog Number: 29958-1-AP	GenBank Accession Number: BC145651	Purification Method: Antigen affinity purification
Size: 600 µg/ml	GeneID (NCBI): 214763	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q8C6L5	
Isotype: IgG	Full Name: RIKEN cDNA E330016A19 gene	
Immunogen Catalog Number: AG31677	Calculated MW: 58 kDa	
	Observed MW: 60 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HT-29 cells, RAW 264.7 cells, Sp2/O cells IHC : human placenta tissue, human lung tissue, mouse lung tissue
Cited Applications: WB, IF	
Species Specificity: Human, Mouse	
Cited Species: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

cGAS (cyclic GMP-AMP synthase) is a cytosolic DNA sensor that serves to mount an immune response against the invasion of microbial pathogens such as viruses. cGAS normally resides as an inactive protein in the cell. Upon binding to DNA, cGAS undergoes a conformational change to an active state and produces the second messenger cyclic GMP-AMP (cGAMP) from ATP and GTP, which is subsequently detected by the cyclic-dinucleotide sensor STING, an ~40 kDa dimeric transmembrane protein at the endoplasmic reticulum (ER). cGAS not only is found in the cytosol but has a multifaceted cellular distribution that involves localization at the cell membrane and in the nucleus (PMID: 32424334). The calculated molecular weight of cGAS is 58 kDa. With post-translational modification, the MW of cGAS will be migrated to 70 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Yupei Jia	38421179	J Virol	WB
Li Jin	38481810	Int J Biol Sci	WB
Tao Yang	38220141	Eur J Pharmacol	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 29958-1-AP (cGAS antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 29958-1-AP (cGAS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).