For Research Use Only

## GPC3 Polyclonal antibody

Catalog Number:30021-1-AP





Basic Information	Catalog Number: 30021-1-AP	GenBank Accession Number: BC035972	Purification Method: Antigen affinity purification					
	Size: 450 µ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG32559	GeneID (NCBI): 2719 UNIPROT ID: P51654 Full Name: glypican 3 Calculated MW: 580 aa, 66 kDa Observed MW: 66 kDa	Recommended Dilutions: WB 1:1000-1:8000 IHC 1:50-1:500					
				Applications	Tested Applications:	Positive C	tive Controls:	
					IHC, WB, ELISA WB : HuH Species Specificity: IHC : mou Human, mouse		7 cells, se liver tissue,	
					Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
				Background Information	Glypicans (GPCs) are a family of glycosylphosphatidylinositol (GPI)-anchored heparan sulfate proteoglycans (HSPGs) that may play a role in the control of cell division and growth regulation. In mammals, there are six GPCs (GPC1 to GPC6), all of which have a similar core-protein size of approx. 60 kDa and the clustering of glycosaminoglycan attachment site near the C-terminus. They are tethered to the cell surface by GPI linkages, which can be cleaved by endogenous phospholipases, thus releasing the protein. Glypican 3 (GPC3) is highly expressed in many tissues during development and plays an important role in the regulation of embryonic growth (PMID: 22467855). Loss-of-function mutations of GPC3 result in the Simpson-Golabi-Behmel overgrowth syndrome (SGBS), and Gpc-3 null mice display developmental overgrowth (PMID: 8589713; 18477453). In hepatocellular carcinoma (HCC), the overexpression of glypican 3 has been demonstrated to be a reliable diagnostic indicator (PMID: 19212669; 22706665). The calculated molecular weight of native glypican 3 is 66 kDa, and glycinate forms o glypican 3 have higher molecular weights than 66 kDa (PMID: 12851874; 16024626; 19574424).			
	expressed in many tissues during (PMID: 22467855). Loss-of-functior (SGBS), and Gpc-3 null mice displa carcinoma (HCC), the overexpress (PMID: 19212669; 22706665). The	development and plays an important n mutations of GPC3 result in the Simp y developmental overgrowth (PMID: ion of glypican 3 has been demonstra calculated molecular weight of native	role in the regulation of embryonic growth oson-Golabi-Behmel overgrowth syndrome 8589713; 18477453). In hepatocellular ted to be a reliable diagnostic indicator a glypican 3 is 66 kDa, and glycinate forms o					

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 30021-1-AP (GPC3 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 30021-1-AP (GPC 3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).