

For Research Use Only

PSMB7 Polyclonal antibody

Catalog Number: 30283-1-AP



Basic Information

Catalog Number: 30283-1-AP	GenBank Accession Number: BC000509	Purification Method: Antigen affinity purification
Size: 550 µg/ml	GeneID (NCBI): 5695	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q99436	
Isotype: IgG	Full Name: proteasome (prosome, macropain) subunit, beta type, 7	
Immunogen Catalog Number: AG32201	Calculated MW: 29.9 kDa	
	Observed MW: 28-30 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, MCF-7 cells, HepG2 cells, mouse skeletal muscle tissue, rat skeletal muscle tissue, mouse kidney tissue, rat kidney tissue

IHC: human colon cancer tissue,

Background Information

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. PSMB7 protein is a member of the proteasome B-type family, also known as the T1B family, and is a 20S core beta subunit in the proteasome. Expression of this catalytic subunit is downregulated by gamma interferon, and proteolytic processing is required to generate a mature subunit. Component of the 20S core proteasome complex are involved in the proteolytic degradation of most intracellular proteins. PSMB7 displays a trypsin-like activity. (PMID:15244466, PMID:27176742, PMID: 8610016)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30283-1-AP (PSMB7 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 30283-1-AP (PSMB7 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 30283-1-AP (PSMB7 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 30283-1-AP (PSMB7 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).