For Research Use Only

ARID1A Polyclonal antibody

Catalog Number:30304-1-AP



Basic Information

Catalog Number: GenBank Accession Number: 30304-1-AP NM_006015
Size: GeneID (NCBI): 400 µ g/ml 8289
Source: UNIPROT ID: Rabbit 014497

Isotype: Full Name:
IgG AT rich interactive domain 1A (SWI-

Immunogen Catalog Number: like)

AG30749 Calculated MW: 242 kDa

Observed MW: 250-260 kDa

Applications

Tested Applications: IHC, IP, WB, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HeLa cells, HepG2 cells, K-562 cells, MCF-7 cells

Purification Method:

WB 1:2000-1:12000

protein lysate

IHC 1:500-1:2000

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

IP: HeLa cells,

IHC: human endometrial cancer tissue,

Background Information

ARID1A, also named as BAF250, BAF250A, C1orf4, OSA1 and SMARCF1, is involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). It binds DNA non-specifically. It is also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene. ARID1A belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The antibody is specific to ARID1A.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 30304-1-AP (ARID1A antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




IP result of anti-ARID1A (IP:30304-1-AP, 4ug; Detection:30304-1-AP 1:3000) with HeLa cells lysate 1280 ug.




Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 30304-1-AP (ARID1A antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 30304-1-AP (ARID1A antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 30304-1-AP (ARID1A antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human endometrial cancer tissue slide using 30304-1-AP (ARID1A antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).