## For Research Use Only

## SLAMF7/CD319 Polyclonal antibody

Catalog Number: 30573-1-AP



**Basic Information** 

Catalog Number: 30573-1-AP

Size:

Isotype:

GenBank Accession Number:

BC027867 GenelD (NCBI):

 400 ug/ml
 57823

 Source:
 UNIPROT ID:

 Rabbit
 Q9NQ25

gG SLAM family member 7

Calculated MW: 335 aa, 37 kDa Observed MW: 66-80 kDa

Full Name:

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:6000 IHC 1:200-1:800

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity:

human

Positive Controls: WB:IM-9 cells,

IHC: human tonsillitis tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

## **Background Information**

Signaling lymphocyte activation molecule (SLAM) family receptors is a group of type I transmembrane receptors belonging to the immunoglobulin superfamily (PMID: 37251372). The SLAM family contains nine members that contain an extracellular segment comprising two or four Ig-like domains (V-like variable and C2-like constant), a transmembrane region and a cytoplasmic tail (PMID: 29662641). They are involved in various physiological and pathological processes, including the regulation of immunological responses and the route of viral entry. SLAMF7, also known as CS1, CRACC or CD319, is expressed on NK cells, CD8+T lymphocytes, B lymphocytes, and mature dendritic cells (PMID: 28861320). It exerts both activating and inhibitory effects depending on the expression status of SAP or EAT-2. SLAMF7 is overexpressed in multiple myeloma and makes it a target for immunotherapy (PMID: 28861320). The native molecular weight of SLAMF7 is 37 kDa and the apparent molecular mass shifts due to glycosylation (PMID: 11698418; 36381324).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 30573-1-AP (SLAMF7/CD319 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 30573-1-AP (SLAMF7 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 30573-1-AP (SLAMF7 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).