For Research Use Only

TIMP1 Polyclonal antibody

Catalog Number: 30755-1-AP



Basic Information

Catalog Number: 30755-1-AP

Size: 800 μ g/ml Source: Rabbit Isotype:

Full Name: TIMP metallopeptidase inhibitor 1 Calculated MW:

> 23 kDa Observed MW: 23-28 kDa

BC000866

7076

P01033

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

Purification Method:

Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF 1:200-1:800

Applications

Tested Applications: WB, IF/ICC, IHC, ELISA Species Specificity: Human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: A2780 cells, IHC: rat testis tissue, IF: HepG2 cells, HeLa cells

Background Information

TIMP1 is a member of the family of matrix metalloproteinase inhibitors, which contains four members (TIMP1, TIMP2, TIMP3, and TIMP4). Tissue inhibitors of metalloproteinases (TIMPs) are multifaceted molecules that exhibit properties beyond their classical proteinase inhibitory function. TIMP1 has several MMP-independent functions such as modulation of angiogenesis, promotion of cell proliferation, and inhibition of apoptosis. TIMP 1 plays importantrole in cell cycle regulation and cancer progression. Recently, clinical studies have shown that the aberrant expression of TIMP1 is associated with an unfavorable prognosis in a series of tumors, such as gastric cancer, papillary thyroid carcinoma, cutaneous melanoma and breast cancer. In pregnancy, TIMP1 plays a regulatory role in the process of implantation, particularly the cytotrophoblast invasion of the uterine endometrium. In pregnancy, TIMP1 plays a regulatory role in the process of implantation, particularly the cytotrophoblast invasion of the uterine endometrium.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




A 2780 cells were subjected to SDS PAGE followed by western blot with 30755-1-AP (TIMP1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded rat testis tissue slide using 30755-1-AP (TIMP1 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded rat testis tissue slide using 30755-1-AP (TIMP1 antibody) at dilution of 1:200 (under 40x lens)



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using TIMP1 antibody (30755-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using TIMP1 antibody (30755-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).