For Research Use Only

Cathepsin C Polyclonal antibody, PBS Only



Purification Method:

Antigen affinity purification

Catalog Number: 30790-1-PBS

Basic Information

Catalog Number:

30790-1-PBS

Size: 1 mg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG32420

52 kDa Observed MW: 100 kDa

NM_001814

UNIPROT ID:

cathepsin C Calculated MW:

P53634 Full Name:

GeneID (NCBI):

GenBank Accession Number:

Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human

Background Information

Applications

Cathepsin C, also known as DPP-I. It is expected to be located in lysosome and highly expressed in lung, kidney and placenta (PMID: 9092576). This gene encodes a member of the peptidase C1 family and lysosomal cysteine proteinase that appears to be a central coordinator for activation of many serine proteinases in cells of the immune system. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate heavy and light chains that form a disulfide-linked dimer. Defects in the encoded protein have been shown to be a cause of Papillon-Lefevre syndrome, an autosomal recessive disorder characterized by palmoplantar keratosis and periodontitis. The calculated molecular weight of Cathepsin C is 51 kDa. The 23-kDa band may be a subunit of the protein monomer, while a 100-kda band is a dimer form (PMID: 26884336, 31557781).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 30790-1-AP (Cathepsin C antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 30790-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded skin cancer slide using 30790-1-AP (Cathepsin C antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 30790-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using Cathepsin C antibody (30790-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 30790-1-PBS in a different storage buffer formulation.