## For Research Use Only

## CD147 Polyclonal antibody

Catalog Number: 30880-1-AP



**Basic Information** 

Catalog Number: 30880-1-AP

GenBank Accession Number: BC009040 Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

350 ug/ml

UNIPROT ID: P35613 WB 1:5000-1:50000 IHC 1:4000-1:16000 IF/ICC 1:200-1:800

Source: Rabbit Isotype:

Full Name:

G basigin (Ok blood group)

Calculated MW: 385 aa, 42 kDa Observed MW: 35-55 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: Jurkat cells, HepG2 cells IHC: human placenta tissue,

IF/ICC: HeLa cells,

## **Background Information**

CD147, also known as Basigin or extracellular matrix metalloproteinase inducer (EMMPRIN), is a transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 7812975). The molecule is composed of an intracellular portion, an extracellular portion and a single transmembrane region. CD147 is expressed on a variety of cell types (e.g., hematopoietic, epithelial, and endothelial cells) and at varying levels (PMID: 32968061). Increased expression of CD147 occurs in many tumors. CD147 is a pleiotropic molecule that plays an important role in fetal, neuronal, lymphocyte and extracellular matrix development (PMID: 17945211). CD147 has been identified as a receptor essential for erythrocyte invasion by Plasmodium falciparum (PMID: 22080952). It has been reported that spike protein of SARS-CoV-2 binds to CD147 on host cells, thereby mediating the viral invasion (Wang, Ke, et al, BioRxiv, 2020).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 30880-1-AP (CD147 antibody) at dilution of 1:25000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 30880-1-AP (CD147 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 30880-1-AP (CD147 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CD147 antibody (30880-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).