### For Research Use Only

# Cullin 5 Polyclonal antibody

Catalog Number: 31183-1-AP



**Basic Information** 

Catalog Number: 31183-1-AP

GeneID (NCBI): Size: 250 ug/ml 8065

**UNIPROT ID:** Source: Rabbit Q93034 Full Name: Isotype: cullin 5 Calculated MW: Immunogen Catalog Number:

AG34620 Observed MW:

80 kDa, 90 kDa

91 kDa

GenBank Accession Number:

BC063306

**Purification Method:** Antigen affinity Purification Recommended Dilutions: WB 1:1000-1:10000

IHC 1:50-1:500

**Applications** 

**Tested Applications:** WB, IHC, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

#### Positive Controls:

WB: LNCaP cells, MCF-7 cells, mouse brain tissue, rat

brain tissue

IHC: mouse brain tissue,

### **Background Information**

The cullin family proteins are scaffold proteins for the RING finger type E3 ligases, participating in the proteolysis through the ubiquitin-proteasome pathway. In humans, there are eight members of cullins, including CUL1, 2, 3, 4A, 4B, 5, 7, and CUL9 with an evolutionarily conserved cullin homology domain at the C-terminus, responsible for their interaction with the RING protein, assembling 8 cullin-based RING dependent E3 ubiquitin ligases to target thousands of substrates for ubiquitination and degradation (PMID: 32334051). Cullin 5 is the key scaffolding protein of Cullin RING ligase-5 (CRL-5). Cullin 5 plays positive roles in regulating cell growth, proliferation and physiological and other processes in the human body (PMID: 32194910). Cullin 5 has been detected in the form of double bands in the results of Western blot, which might be caused by neddylation (PMID: 36449095; 31294695).

## Storage


#### Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 31183-1-AP (Cullin 5 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 31183-1-AP (Cullin 5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 31183-1-AP (Cullin 5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 31183-1-AP (Cullin 5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).