For Research Use Only

## HAPLN1 Polyclonal antibody

Catalog Number:31646-1-AP




Basic Information	Catalog Number: 31646-1-AP	GenBank Accession Number: BC057808	Purification Method: Antigen affinity Purification				
	<mark>Size:</mark> 650 μg/ml	GenelD (NCBI): 1404	Recommended Dilutions: WB 1:500-1:1000				
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG35652	UNIPROT ID: P10915					
		Full Name: hyaluronan and proteoglycan link protein 1 Calculated MW: 40 kDa Observed MW: 40 kDa					
				Applications	Tested Applications: WB, ELISA Species Specificity: mouse, rat	Positive Controls:	
						WB : mous	se brain tissue, rat brain tissue
Background Information	cytoplasm and extracellular, and molecular weight of HAPLN1 is 40 aggregates and thus increasing co factor, up regulating aggrecan and	HAPLN1 (hyaluronan and proteoglycan link protein 1), also known as CRTL1. It is expected to be located in the cytoplasm and extracellular, and the protein is mainly expressed in placenta and small intestine. The calculated molecular weight of HAPLN1 is 40 kDa. HAPLN1 binds to aggrecan on the hyaluronic chain, stabilizing the aggregates and thus increasing compression resistance and shock absorption. HAPLN1 also functions as a growth factor, up regulating aggrecan and type II collagen synthesis in cartilage. It seems to be an essential part of cartilage homeostasis and may also contribute to homeostasis in the disc tissue (PMID: 23537453).					
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	1 50% glycerol pH 7.3.					

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31646-1-AP (HAPLN1 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.