For Research Use Only

VE-cadherin/CD144 Polyclonal antibody

Catalog Number: 31772-1-AP



Basic Information

Catalog Number: 31772-1-AP

Size: 350 ug/ml Source: Rabbit

Isotype:

GenBank Accession Number:

NM_001795.5 GeneID (NCBI): 1003 **UNIPROT ID:**

P33151-1 Full Name:

cadherin 5, type 2 (vascular endothelium)

Calculated MW: 88 kDa

Observed MW: 90-100 kDa, 130-140 kDa **Purification Method:** Antigen affinity Purification Recommended Dilutions: WB 1:2000-1:16000 IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IF/ICC, ELISA Species Specificity:

Positive Controls:

WB: HUVEC cells, human placenta tissue

IF/ICC: HUVEC cells,

Background Information

 $Cadherins\ are\ a\ family\ of\ transmembrane\ gly coproteins\ that\ mediate\ calcium-dependent\ cell-cell\ adhesion\ and$ play an important role in the maintenance of normal tissue architecture. Vascular endothelial cadherin (VEcadherin), also known as Cadherin-5 (CDH5) or CD144, is a member of the type II classical cadherin family of cell adhesion proteins (PMID: 21269602). VE-cadherin is expressed specifically in endothelial cells and mediates homophilic adhesion in the vascular endothelium (PMID: 1522121; 8555485; 21269602). VE-cadherin plays a role in the organization of lateral endothelial junctions and in the control of permeability properties of vascular endothelium (PMID: 1522121). VE-cadherin has also been shown to be required for angiogenesis (PMID: 16473763; 18162609). The calculated molecular weight of VE-cadherin is 88 kDa and the apparent molecular weight of 120-140 kDa is higher due to post-translational glycosylation and phosphorylation (PMID: 10460833; 29894844). Fulllength VE-cadherin can be proteolytically cleaved to generate a fragment of 90-100 kDa (PMID: 9786462; 22064597).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31772-1-AP (VEcadherin/CD144 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using VE-cadherin/CD144 antibody (31772-1-AP) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).