For Research Use Only

CPN1 Polyclonal antibody Catalog Number: 31940-1-AP


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 31940-1-AP	GenBank Accession Number: BC027897	Purification Method: Antigen affinity Purification
	Size: 550 ug/ml	GenelD (NCBI): 1369	Recommended Dilutions: WB 1:1000-1:6000
	Source: Rabbit	UNIPROT ID: P15169	
	lsotype: IgG		
	Immunogen Catalog Number: AG34806	Calculated MW: 458 aa, 52 kDa	
		Observed MW: 40~50 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, ELISA Species Specificity: human	WB : human plasma,	
	nunan		
Background Information	Human carboxypeptidase N (CPN) an extracellular glycoprotein sym vasoactive peptide hormones, gro Normally, CPN circulates in blood flanked by a 48 to 55 kDa catalyti family members in that it contain	thesized in the liver and secreted into wth factors and cytokines by specifica plasma as a hetero-tetramer consistir c (CPN1) domain. The C terminal regio	the blood, where it controls the activity of Illy removing C-terminal basic residues. ng of two 83 kDa (CPN2) domains each on of the CPN1 subunit differs from all other s at these sites apparently happens either
Background Informatio	Human carboxypeptidase N (CPN) an extracellular glycoprotein sym vasoactive peptide hormones, gro Normally, CPN circulates in blood flanked by a 48 to 55 kDa catalyti family members in that it contain	thesized in the liver and secreted into with factors and cytokines by specifica plasma as a hetero-tetramer consistir c (CPN1) domain. The C terminal regis s numerous basic residues. Proteolysis ng processing and secretion from the l after shipment.	Illy removing C-terminal basic residues. ng of two 83 kDa (CPN2) domains each on of the CPN1 subunit differs from all other s at these sites apparently happens either

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 31940-1-AP (CPN1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.